

524 Rec'd PCT/PTO 2 8 SEP 1999

FORM PTO-1390 (REV 10-95)		U. S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER PC10030A	
TRANSMITTAL LETTER TO THE UNITED STATES PATENT AND TRADEMARK OFFICE DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.5) Not yet assigned 09/402010	
INTERNATIONAL APPLICATION NO. PCT/IB98/01813		INTERNATIONAL FILING DATE November 13, 1998 (11.13.1998)		PRIORITY DATE CLAIMED December 31, 1997 (12.31.1997)	
TITLE OF INVENTION ARYL FUSED AZAPOLYCYCLIC COMPOUNDS					
APPLICANT(S) FOR DO/EO/US Jotham Wadsworth COE and Paige Roanne Palmer BROOKS					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is the FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is the SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 					
Items 11. To 16. Below concern other documents(s) or information included:					
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input type="checkbox"/> A change of power of attorney and/or address letter. <input type="checkbox"/> Other items or information: 					

EXPRESS MAIL NO. EM 984852791

TRANSMITTAL LETTER UNDER 35 U.S.C. 371 PTO 1390, 3/99

514 Rec'd PCT/PTO 2 8 SEP 1999

U.S. APPLICATION NO. (If known, see 37 CFR 1.5) Not yet assigned 09/402010		INTERNATIONAL APPLICATION NO. PCT/IB98/01813		ATTORNEY'S DOCKET NUMBER PC10030A	
17. <input checked="" type="checkbox"/> The following fees are submitted BASIC NATIONAL FEE (37 CFR 1.492 (a)(1)-(5)): <input checked="" type="checkbox"/> Search Report has been prepared by the EPO or JPO\$840.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37CFR 1.482)\$670.00 <input type="checkbox"/> No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2))\$760.00 <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO\$970.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)\$ 96.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	- 20 =		X \$ 18.00	\$	
Independent Claims	- 3 =		X \$ 78.00	\$78	
MULTIPLE DEPENDENT CLAIM(s) (if applicable)			+ \$260.00	\$	
TOTAL OF ABOVE CALCULATIONS =				\$918	
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed. (Note: 37 CFR 1.9, 1.27, 1.28)				\$	
SUBTOTAL =				\$918	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$918	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				\$	
TOTAL FEES ENCLOSED =				\$918	
				Amount to be: Refunded \$ Charged \$	
a. <input type="checkbox"/> A check in the amount of \$ _____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. 16-1445 in the amount of \$ <u>918</u> to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No.16-1445. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Paul H. Ginsburg Pfizer Inc 235 East 42nd Street New York, NY 10017-5755					
				<i>Karen DeBenedictis</i> Signature Karen DeBenedictis Name 32,977 Registration Number	

EXPRESS MAIL NO. EM 484852791

TRANSMITTAL LETTER UNDER 35 U.S.C. 371 PTO 1390, 3/99

5

ARYL FUSED AZAPOLYCYCLIC COMPOUNDSBackground of the Invention


This invention relates to aryl fused azapolycyclic compounds, as defined more specifically by formula I below. Compounds of formula I bind to neuronal nicotinic acetylcholine specific receptor sites and are useful in modulating cholinergic function. Such compounds are useful in the treatment of inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS), cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), obsessive-compulsive disorder, psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia, dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome.

The compounds of this invention may also be used in combination with an antidepressant such as, for example, a tricyclic antidepressant or a serotonin reuptake inhibiting antidepressant (SRI), in order to treat both the cognitive decline and depression associated with AD, PD, stroke, Huntington's Chorea or traumatic brain injury (TBI); in combination with muscarinic agonists in order to stimulate both central muscarinic and nicotinic receptors for the treatment, for example, of ALS, cognitive dysfunction, age related cognitive decline, AD, PD, stroke, Huntington's Chorea and TBI, in combination with neurotrophic factors such as NGF in order to maximize cholinergic enhancement for the treatment, for example, of ALS, cognitive dysfunction, age related cognitive decline, AD, PD stroke, Huntington's Chorea and TBI; or in combination with agents that slow or arrest AD such as cognition enhancers, amyloid aggregation inhibitors, secretase inhibitors, tau kinase inhibitors, neuronal antiinflammatory agents and estrogen-like therapy

5 Other compounds that bind to neuronal nicotinic receptor sites are referred to in United States Patent Application 08/963,852, which was filed on November 4, 1997. The foregoing application is owned in common with the present application, and is incorporated herein by reference in its entirety.

Summary of the Invention

10 This invention relates to aryl fused azapolycyclic compounds of the formula



R¹ is hydrogen, (C₁-C₆)alkyl, unconjugated (C₃-C₆)alkenyl, benzyl, XC(=O)R¹³ or -CH₂CH₂-O-(C₁-C₄)alkyl;

R² and R³ are selected, independently, from hydrogen, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, hydroxy, nitro, amino, halo, cyano, -SO_q(C₁-C₆)alkyl wherein q is zero, one or two, (C₁-C₆)alkylamino-, [(C₁-C₆)alkyl]₂amino-, -CO₂R⁴, -CONR⁵R⁶, -SO₂NR⁷R⁸, -C(=O)R¹³, -XC(=O)R¹³, aryl-(C₀-C₃)alkyl- or aryl-(C₀-C₃)alkyl-O-, wherein said aryl is selected from phenyl and naphthyl, heteroaryl-(C₀-C₃)alkyl- or heteroaryl-(C₀-C₃)alkyl-O-, wherein said heteroaryl is selected from five to seven membered aromatic rings containing from one to four heteroatoms selected from oxygen, nitrogen and sulfur, and X²(C₀-C₆)alkoxy-(C₀-C₆)alkyl-, wherein X² is absent or X² is (C₁-C₆)alkylamino- or [(C₁-C₆)alkyl]₂amino-, and wherein the (C₀-C₆)alkoxy-(C₀-C₆)alkyl- moiety of said X²(C₀-C₆)alkoxy-(C₀-C₆)alkyl- contains at least one carbon atom, and wherein from one to three of the carbon atoms of said (C₀-C₆)alkoxy-(C₀-C₆)alkyl- moiety may optionally be replaced by an oxygen, nitrogen or sulfur atom, with the proviso that any two such heteroatoms must be separated by at least two carbon atoms, and wherein any of the alkyl moieties of said (C₀-C₆)alkoxy-(C₀-C₆)alkyl- may be optionally substituted with from two to seven fluorine atoms, and wherein one of the carbon atoms of each of the alkyl moieties of said aryl-(C₀-C₃)alkyl- and said heteroaryl-(C₀-C₃)alkyl- may optionally be replaced by an oxygen, nitrogen or sulfur atom, and wherein each of the foregoing aryl and heteroaryl groups may optionally be substituted with one or more substituents, preferably from zero to two substituents, independently selected from (C₁-C₆)alkyl optionally substituted with from one to seven fluorine atoms, (C₁-C₆)alkoxy optionally substituted with from two to seven fluorine atoms, halo (e.g., chloro fluoro, bromo or iodo), (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, hydroxy, nitro, cyano, amino, (C₁-

- 5 C₆)alkylamino-, [(C₁-C₆) alkyl]₂amino-, -CO₂R⁴, -CONR⁵R⁶, -SO₂NR⁷R⁸, -C(=O)R¹³ and -XC(=O)R¹³,







or R² and R³, together with the carbons to which they are attached, form a four to seven membered monocyclic, or a ten to fourteen membered bicyclic, carbocyclic ring that can be saturated or unsaturated, wherein from one to three of the nonfused carbon atoms of said
 10 monocyclic rings, and from one to five of the carbon atoms of said bicyclic rings that are not part of the benzo ring shown in formula I, may optionally and independently be replaced by a nitrogen, oxygen or sulfur, and wherein said monocyclic and bicyclic rings may optionally be substituted with one or more substituents, preferably from zero to two substituents for the monocyclic rings and from zero to three substituents for the bicyclic rings, that are selected,
 15 independently, from (C₀-C₆)alkoxy-(C₀-C₆)alkyl-, wherein the total number of carbon atoms does not exceed six and wherein any of the alkyl moieties may optionally be substituted with from one to seven fluorine atoms; nitro, oxo, cyano, halo, (C₂-C₆)alkenyl, (C₂-C₆)alkynyl, hydroxy, amino, (C₁-C₆)alkylamino-, [(C₁-C₆)alkyl]₂amino-, -CO₂R⁴, -CONR⁵R⁶, -SO₂NR⁷R⁸, -C(=O)R¹³, and -XC(=O)R¹³,

20 each R⁴, R⁵, R⁶, R⁷, R⁸ and R¹³ is selected, independently, from hydrogen and (C₁-C₆) alkyl, or R⁵ and R⁶, or R⁷ and R⁸ together with the nitrogen to which they are attached, form a pyrrolidine, piperidine, morpholine, azetidine, piperazine, -N-(C₁-C₆)alkylpiperazine or thiomorpholine ring, or a thiomorpholine ring wherein the ring sulfur is replaced with a sulfoxide or sulfone; and

25 each X is, independently, (C₁-C₆)alkylene,
 with the proviso that. (a) at least one of R¹, R² and R³ must be the other than hydrogen, and (b) when R² and R³ are hydrogen, R¹ cannot be methyl or hydrogen;
 and the pharmaceutically acceptable salts of such compounds.

Examples of heteroaryl groups that each of R² and R³ can be are the following:

30 thienyl, oxazolyl, isoxazolyl, pyridyl, pyrimidyl, thiazolyl, tetrazolyl, isothiazolyl, triazolyl, imidazolyl, tetrazolyl, pyrrolyl and the following groups








5

wherein one of R^9 and R^{18} is hydrogen or (C_1-C_6) alkyl, and the other is a bond to the benzo ring of formula I.

Examples of compounds of this invention are compounds of the formula I, and their pharmaceutically acceptable salts, wherein R^2 and R^3 , together with the benzo ring of formula I,


10 form a bicyclic ring system selected from the following:













wherein R^{10} and R^{17} are selected, independently, from (C_0-C_6) alkoxy- (C_0-C_6) alkyl- wherein the total number of carbon atoms does not exceed six and wherein any of the alkyl moieties may optionally be substituted with from one to seven fluorine atoms; nitro, cyano, halo, amino, (C_1-C_6) alkylamino-, $[(C_1-C_6) \text{ alkyl}]_2$ amino-, $-CO_2R^4$, $-CONR^5R^6$, $-SO_2NR^7R^8$, $-C(=O)R^{13}$, $-XC(=O)R^{13}$, phenyl and monocyclic heteroaryl wherein said heteroaryl is defined as R^2 and R^3 are defined in the definition of compounds of the formula I above;

Other embodiments of this invention relate to compounds of the formula I, and their pharmaceutically acceptable salts, wherein R^2 and R^3 , together with the benzo ring of formula I, form a bicyclic or tricyclic ring system selected from the following:

20



5



wherein R^{10} and R^{17} are defined as above and m is zero, one or two, and wherein one of the carbon atoms of ring A can optionally be replaced with oxygen or $-N(C_1-C_6)\text{alkyl}$.

Other embodiments of this invention relate to compounds of the formula I, and their pharmaceutically acceptable salts, wherein neither R^2 nor R^3 is attached to the benzo ring of formula I via an oxygen atom.

Other embodiments of this invention relate to compounds of the formula I wherein R^1 is not methyl

Examples of specific compounds of the formula I are the following:

6-methyl-5,7-dioxo-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;

- 5 6-methyl-5-oxo-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
5,7-dimethyl-6-oxo-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
5,7-dioxo-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene
- 10 hydrochloride;
5-oxo-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
6-oxo-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
4,5-difluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride;
5-fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-carbonitrile hydrochloride;
- 15 4-ethynyl-5-fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride;
5-ethynyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-carbonitrile hydrochloride;
5-chloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-carbonitrile hydrochloride;
4-ethynyl-5-chloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride;
5-oxa-7-methyl-6-oxo-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene
- 20 hydrochloride;
4-fluoro-5-trifluoromethyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride;
4-chloro-5-trifluoromethyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride;
- 25 5-trifluoromethyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-carbonitrile hydrochloride;
4-ethynyl-5-trifluoromethyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride,
- 30 6-methyl-5-thia-5-dioxo-6,13-Diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
7-dimethylamino-5-thia-5-dioxo-6,13-Diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,8-triene hydrochloride;
6,7-dioxo-5,8,14-triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,9-triene hydrochloride; and
- 35 5,8-dimethyl-6,7-dioxo-5,8,14-triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,9-triene hydrochloride

This invention also relates to compounds of the formula



10

15

20

20

25

25

30

The compounds of formula I may have optical centers and therefore may occur in different enantiomeric configurations. The invention includes all enantiomers, diastereomers, and

5 other stereoisomers of such compounds of formula I, as well as racemic and other mixtures thereof.

The present invention also relates to all radiolabelled forms of the compounds of the formulae I. Preferred radiolabelled compounds of formula I are those wherein the radiolabels are selected from as ^3H , ^{11}C , ^{14}C , ^{18}F , ^{123}I and ^{125}I . Such radiolabelled compounds are useful as
10 research and diagnostic tools in metabolism pharmacokinetics studies and in binding assays in both animals and man.

The present invention also relates to a pharmaceutical composition for use in reducing nicotine addiction or aiding in the cessation or lessening of tobacco use in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically
15 acceptable salt thereof, that is effective in reducing nicotine addiction or aiding in the cessation or lessening of tobacco use and a pharmaceutically acceptable carrier.


The present invention also relates to a method for reducing nicotine addiction or aiding in the cessation or lessening of tobacco use in a mammal, including a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically
20 acceptable salt thereof, that is effective in reducing nicotine addiction or aiding in the cessation or lessening of tobacco use.

The present invention also relates to a method of treating a disorder or condition selected from inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia,
25 chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS), cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco
30 products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia, dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal,
35 comprising administering to a mammal in need of such treatment an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, that is effective in treating such disorder or condition.

09402010-092009

5 The present invention also relates to a pharmaceutical composition for treating a disorder or condition selected from inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral
10 sclerosis (ALS), cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive
15 dyskinesia, hyperkinesia, dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.


20 The present invention also relates to a method for reducing nicotine addiction or aiding in the cessation or lessening of tobacco use in a mammal, comprising administering to said mammal an amount of a compound comprising an amount of a compound of the formula



or a pharmaceutically acceptable salt thereof, that is effective in reducing nicotine
25 addiction or aiding in the cessation or lessening of tobacco use

 The present invention also relates to a method for treating a disorder or condition selected from inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS),
30 cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia,
35 dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including

- 5 petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal, comprising administering to a mammal in need of such treatment an amount of a compound of the formula



- 10 or a pharmaceutically acceptable salt thereof, that is effective in treating such disorder or condition.


- This invention also relates to the pharmaceutically acceptable acid addition salts of the compounds of formula I. Examples of pharmaceutically acceptable acid addition salts of the compounds of formula I are the salts of hydrochloric acid, p-toluenesulfonic acid, fumaric acid, 15 citric acid, succinic acid, salicylic acid, oxalic acid, hydrobromic acid, phosphoric acid, methanesulfonic acid, tartaric acid, malate, di-p-toluoyl tartaric acid, and mandelic acid.

Detailed Description of the Invention

Except where otherwise stated, R¹ through R¹⁸, m and P, and structural formula I in the reaction schemes and discussion that follow are defined as above.


5

Scheme 1




5

Scheme 2




5

Scheme 2 continued





5

Scheme 3



5


Scheme 4



653250 " 07020450

5


Scheme 5



09402010, 052334

5



Scheme 6



034022010 032239
652250 01020450

5


Scheme 6 continued



094022010 0928379 658250 " 07020460

5

Scheme 7




IG: (R^2 and R^3 form ring A)

III: (ring A = absent)

5


Scheme 8



04022010 092839


5

Scheme 9



5

Scheme 10



5 Scheme 1-10 illustrate methods of synthesizing compounds of the formula I .

Referring to Scheme 1, the starting material of formula III is reacted with trifluoroacetic anhydride, in the presence of pyridine, to form the compound of formula IV. This reaction is typically conducted in methylene chloride at a temperature from about 0°C to about room temperature.


10 The compound of formula IV is then converted into the dinitro derivative of formula IIA by the following process. The compound of the formula IV is added to a mixture of 4 or more equivalents of trifluoromethanesulfonic acid ($\text{CF}_3\text{SO}_2\text{OH}$) and 2 to 3 equivalents of nitric acid, in a chlorinated hydrocarbon solvent such as chloroform, dichloroethane (DCE) or methylene chloride. The resulting mixture is allowed to react for about 5 to 24 hours. Both of the foregoing
15 reactions are generally conducted at a temperature ranging from about -78°C to about 0°C for about 2 hours, and then allowed to warm to room temperature for the remaining time.

Reduction of the compound of formula IIA, using methods well known to those of skill in the art, yields the compound of formula IIB. This reduction can be accomplished, for example, using hydrogen and a palladium catalyst such as palladium hydroxide and running the reaction
20 in methanol at about room temperature.

Referring to Scheme 2, the compound of formula IIA is converted into the corresponding compound wherein the trifluoroacetyl protecting group is replaced by a t-Boc protecting group (VIA) by reacting it first with an alkali metal or alkaline earth metal (or ammonium) hydroxide or carbonate, and then reacting the isolated product from the foregoing reaction with di-t-
25 butyldicarbonate. The reaction with the alkali or alkaline earth metal (or ammonium) hydroxide or carbonate is generally carried out in an aqueous alcohol, dioxane or tetrahydrofuran (THF) at a temperature from about room temperature to about 70°C, preferably at about 70°C, for about one to about 24 hours. The reaction of the isolated, unprotected amine or an acid addition salt of such amine, from the above reaction with di-t-butyldicarbonate is preferably carried out in a
30 solvent such as THF, dioxane or methylene chloride at a temperature from about 0°C to about room temperature. This reaction may or may not be conducted in the presence of a base. When the reactant is a salt of the amine, use of a base is preferred. The resulting compound of formula VIA can be converted into the corresponding diamino derivative of formula VIB using the procedure described above for converting the dinitro compound of formula IIA into the
35 corresponding diamino compound of formula IIB.

The conversion of the compound of formula VIB into the desired compound of the formula VII can be accomplished by reacting the compound of formula VIB with a compound of the formula

00402010 09239
1553250 07020460



5 wherein R^{10} is hydrogen, $(\text{C}_1\text{-C}_6)$ alkyl optionally substituted with from one to seven fluorine atoms, aryl- $(\text{C}_0\text{-C}_3)$ alkyl wherein said aryl is selected from phenyl and naphthyl, or heteroaryl- $(\text{C}_0\text{-C}_3)$ alkyl wherein said heteroaryl is selected from five to seven membered aromatic rings containing from one to four heteratoms selected from oxygen, nitrogen and sulfur, and wherein
 10 each of the foregoing aryl and heteroryl groups may optionally be substituted with one or more substituents, preferably from zero to two substituents, independently selected from $(\text{C}_1\text{-C}_6)$ alkyl optionally substituted with from one to seven fluorine atoms, $(\text{C}_1\text{-C}_6)$ alkoxy optionally substituted with from one to seven fluorine atoms and cyano. The preferred solvent for this reaction is a 10:1 mixture of ethanol:acetic acid. The reaction temperature can range from about 40°C to
 15 about 100°C . It is preferably about 60°C . Other appropriate solvents include acetic acid, ethanol and isopropanol.


Alternate methods of preparing compounds of the formula VII the compound of formula VIB are described by Segelstein et al., *Tetrahedron Lett.*, 1993, 34, 1897.

20 Removal of the t-Boc protecting group from the compound of formula VII yields corresponding compound of formula IA. The protecting group can be removed using methods well known to those of skill in the art. For example, the compound of formula VII can be treated with an anhydrous acid such as hydrochloric acid, hydrobromic acid, methanesulfonic acid, or trifluoroacetic acid, preferably hydrochloric acid in ethyl acetate, at a temperature from about 0°C to about 100°C , preferably from about room temperature to about 70°C , for about
 25 one to 24 hours.

The compound of formula VII can be converted into the corresponding compound of formula IB by reacting it with a compound of the formula R^{17}Z , wherein R^{17} is defined as R^{10} is defined above, and Z is a leaving group such as a halo or sulfonate (e.g., chloro, bromo, mesylate or tosylate), in the presence of a base such as an alkali metal hydride, hydroxide or
 30 carbonate, preferably potassium hydroxide, in a polar solvent such as water, dimethylsulfoxide (DMSO), THF or DMF, preferably a mixture of DMSO and water, and then removing the protecting group as described above. The reaction with R^{17}Z is generally carried out at a temperature from about room temperature to about 100°C , preferably at about 50°C , for about five hours.

5 Scheme 3 illustrates an alternate method of preparing compounds of the formula IB from the compound of formula VIA. This method is the preferred method of making compounds of the formula IB wherein R¹⁷ is a bulky group such as an aryl or heteroaryl containing group, or when R¹⁷ can not be attached, as illustrated in Scheme 2, by alkylation or aryl substitution methods. Referring to Scheme 3, the compound of formula VIA is reacted
10 with the appropriate compound of formula R¹⁷NH₂ in a polar solvent such as THF, DMF or DMSO, preferably THF, at a temperature from about room temperature to about 100°C, preferably at the reflux temperature, for about four to eighteen hours. The resulting compound of formula XXIII is then converted into the corresponding compound of the formula XXIV by reducing the nitro group to an amino group using methods well known to those of skill in the art. Such methods are referred to above for the conversion of the compounds of the formula
15 IIA into a compound of the formula IIB in Scheme 1, and exemplified in experimental Examples 12B and 18B. Closure of the imidazole ring to form the corresponding compound of formula XXV can then be accomplished by reacting the compound of formula XXIV from the above reaction with a compound of the formula


20



wherein R¹⁰ is defined as above, as described above for converting compounds of the formula VIB into those of the formula VII


25 Removal of the protecting group from the compound of formula XXV yields the corresponding compound of formula IB. This can be accomplished using methods well known in the art, for example, as described above for forming compounds of the formula IA from the corresponding compounds of the formula VII

Scheme 4 illustrates a method of preparing compounds of the formula IC, wherein R¹⁰ and R¹⁷ are as defined above. Referring to Scheme 4, the compound of formula VIB is
30 reacted with a compound of the formula



5 (sodium bisulfite ethane dione addition adduct) in water or another polar solvent such as THF, DMF or DMSO, preferably a mixture of water and a water miscible solvent such as THF, for about one to four hours. The reaction temperature can range from about 40°C to about 100°C, and is preferably at about the reflux temperature.

10 Alternatively, the compound of formula VIB can be reacted with a compound of the formula



15 (double condensation reaction) in a polar solvent such as THF, water, or acetic acid, preferably a mixture of water and THF. This reaction is typically carried out at a temperature from about 40°C to about 100°C, preferably at the reflux temperature, for about two to four hours.

The desired quinoxaline of formula IC can then be formed by deprotecting the compound formed in either of the foregoing reactions, using the method described above for converting a compound of the formula VII into one of the formula IA

20 Scheme 5 illustrates a method of preparing compounds of the formula I wherein R² and R³, together with the benzo ring to which they are attached, form a benzoxazole ring system. Such a compound, wherein R¹ is hydrogen, is depicted in Scheme 5 as chemical formula IE. Referring to Scheme 5, the compound of formula XXII, wherein Y is nitro, halo, trifluoromethanesulfonate or a diazonium salt, is reacted with potassium acetate or another alkali or alkaline earth metal carboxylate in a solvent such as dimethylsulfoxide (DMSO), DMF or
25 acetonitrile, preferably DMSO. This reaction is generally allowed to run for about 12-24 hours. Appropriate reaction temperatures range from about 70°C to about 140°C. Approximately 100°C is preferred.

The above reaction yields the compound of formula VIII, which can then be converted into the desired compound having formula IE by the following procedure. First, the compound of
30 formula VIII is reduced by reaction with hydrogen and a palladium or platinum catalyst such as palladium hydroxide in methanol at a temperature from about 0°C to about 70°C, preferably at about room temperature, to form the corresponding amino derivative. The product of this reaction is then reacted with an acid chloride of the formula R¹⁰COCl or an acid anhydride of the formula (R¹⁰CO)₂O wherein R¹⁰ is (C₁-C₆)alkyl, or a compound of the formula R¹⁰C(OC₂H₅)₃, in
35 an appropriate inert solvent such as decalin, chlorobenzene or xylenes. A mixture of xylenes is

5 preferred. This reaction is typically conducted at a temperature from about 120-150°C,
preferably at about 140°C. When $R^{10}COCl$ is used as a reactant, it is preferable to add a
stoichiometric amount of triethylamine (TEA) or another organic tertiary amine base and a
catalytic amount of pyridinium p-toluenesulfonic acid or pyridinium p-toluenesulfonate (PPTs) to
the reaction mixture. When $R^{10}C(OC_2H_5)_3$ is used as a reactant, it is preferable to add a catalytic
10 amount of PPTs to the reaction mixture


Removal of the trifluoroacetyl nitrogen protecting group yields the desired compound of
the formula IE. This can be accomplished using methods well known to those of skill in the art,
for example, reacting the protected compound with a lower alkanol and an aqueous alkali or
alkaline earth metal (or ammonium) hydroxide or carbonate, aqueous sodium carbonate, at a
15 temperature from about 50°C to about 100°C, preferably at about 70°C, for about two to six
hours.

Scheme 6 illustrates the preparation of compounds of the formula I wherein R^1 is
hydrogen and R^2 and R^3 , together with the benzo ring to which they are attached, form a
benzothiazole ring system. Referring to Scheme 6, the compound of formula III is reacted with
20 trifluoroacetic anhydride to form the corresponding compound wherein the ring nitrogen is
protected by a trifluoroacetyl group, and the resulting nitrogen protected compound is then
reacted with two equivalents of trifluoromethanesulfonic anhydride and one equivalent of nitric
acid to form the corresponding compound of formula IX, wherein there is a single nitro
substituent on the benzo ring. The reaction with trifluoroacetic acid is typically conducted in the
25 presence of pyridine. Both of the above reactions are typically conducted in a reaction inert
solvent such as a chlorinated hydrocarbon solvent, preferably methylene chloride, at a
temperature from about 0°C to about room temperature, preferably at about room temperature

The above transformation can also be accomplished using other nitration methods
known to those skill in the art

30 Reduction of the nitro group to an amine group can be accomplished as described
above to provide a compound of the formula IX'.

The compound of formula IX' is then reacted with a carboxylic acid halide or anhydride
of the formula $R^{10}COX$ or $(R^{10}CO)_2O$, wherein X is halo and R^{10} is hydrogen or (C_1-C_6) alkyl, and
pyridine, TEA or another tertiary amine base, to form a compound of the formula X, which can
35 then be converted to the desired compound having formula XI by reacting it with Lawesson's
reagent, which is depicted below



5

The reaction with $R^{10}COX$, wherein X is halo, or $(R^{10}CO)_2O$ is generally carried out at a temperature from about $0^{\circ}C$ to about room temperature, preferably at about room temperature. The reaction with Lawesson's reagent is generally carried out in a reaction inert solvent such as benzene or toluene, preferably toluene, at a temperature from about room temperature to about the reflux temperature of the reaction mixture, preferably at about the reflux temperature.

10

Closure to the benzothiazole ring and nitrogen deprotection to form the desired compound of formula IF can be accomplished by reacting the compound of formula XI with potassium ferricyanide and sodium hydroxide in a mixture of water and methanol ($NaOH/H_2O/CH_3OH$), at a temperature from about $50^{\circ}C$ to about $70^{\circ}C$, preferably at about $60^{\circ}C$ for about 1.5 hours.

15

Scheme 7 illustrates a method of preparing the compound of formula III, which is used as the starting material for the process of Scheme 1, or a compound of the formula IG, wherein R^2 and R^3 form a ring (labeled "A" in the Scheme), as defined above in the definition of compounds of the formula I. Referring to Scheme 7, the compound of formula XII, wherein X^1 and X^2 are selected, independently, from chloro, fluoro, bromo and iodo, but where at least one of X^1 and X^2 is Br- or I-, reacted with cyclopentadiene, in the presence of magnesium metal, in a THF, dioxane or other ethereal solvent, at a temperature from about $40^{\circ}C$ to about $100^{\circ}C$, preferably at about the reflux temperature, to form a compound of the formula XIII

20

25 Reaction of the resulting compound of formula XIII with N-methylmorpholine-N-oxide (NMO) and osmium tetroxide in acetone at about room temperature yields the corresponding compound of the formula XIII A.


The compound having formula XIII A is then converted into the corresponding compound of formula XIV using the following procedure. First, the compound of formula XIII A is reacted with sodium periodate in a mixture of a chlorinated hydrocarbon, preferably dichloroethane (DCE), and water, or with lead tetraacetate in a chlorinated hydrocarbon solvent, at a temperature from about $0^{\circ}C$ to about room temperature, to generate a dialdehyde or glycal intermediate. The product of this reaction is then reacted with benzylamine and

30

5 sodium triacetoxyborohydride in a chlorinated hydrocarbon solvent at a temperature from about 0°C to about room temperature, preferably at about room temperature, to form the desired compound of formula XIV. Removal of the benzyl group from the compound of formula XIV yields the compound of formula III (when ring A is absent) or IG, (when ring A is present). This can be accomplished using methods well known to those of skill in the art, for
 10 example, optionally reacting the free base with one equivalent of acid, e.g., hydrochloric acid, (to form the corresponding acid addition salt), followed by hydrogen and palladium hydroxide in methanol at about room temperature.

In the reductive animation step described above and throughout this document, alternatives to benzyl amine, such as ammonia, hydroxylamine, alkoxy amines, methyl amine,
 15 allyl amine, and substituted benzyl amines (e.g., diphenylmethyl amine and 2- and 4-alkoxy substituted benzyl amines) can also be used. They can be used as free bases, or as their salts, preferably their acetate salts, and can be subsequently removed by methods described for each by T W Greene and G M. Wuts, "Protective Groups in Organic Synthesis", 1991, John Wiley & Sons, New York NY

20 The procedure of Scheme 7 can also be used to prepare compounds of the formula I wherein R² and R³ do not form a ring and are not both hydrogen, by replacing the starting material of formula XII with the appropriate compound having the formula



Scheme 8, 9 and 10 illustrate methods of preparing compounds of the formula I
 25 wherein R¹ is hydrogen, and R² and R³ represent a variety of different substituents, as defined above, but do not form a ring.

Scheme 8 illustrates a variation of the process shown in Scheme 7, which can be used to make a compound identical to that of formula III except that the benzo ring is substituted with a fluoro group or an alkoxy group (R¹⁸ in Scheme 8). This compound is depicted in Scheme 8
 30 as chemical structure 1H. Referring to Scheme 8, where, for example, R¹⁸ is F, 1,3-difluorobenzene is reacted with a strong base such as an alkali metal dialkylamine or an alkali metal alkyl (or aryl) in an ethereal solvent such as ethyl ether or THF, at a temperature below -50°C, followed by quenching with iodine or N-iodosuccinamide, to form 1,3-difluoro-2-iodobenzene. The compound 1,3-difluoro-2-iodobenzene (structural formula XVI in Scheme 8)
 35 is then converted into the compound of formula IH by a series of reactions (represented in

5 Scheme 8 as ,XVI→XVII→XVIII→XIX→IH) that are analogous to the series of reactions described above and illustrated in Scheme 7 for converting compounds of the formula XIII into those of the formula IG or III. Conversion of the compound of formula XVI into the compound of formula XVII can also be accomplished by treating a mixture of the compound of formula XVI and cyclopentadiene with an alkyl lithium reagent, preferably n-butyl lithium, in an inert
10 hydrocarbon solvent such as petroleum ether or methyl cyclohexane, at a temperature from about -20°C to about room temperature, preferably at about 0°C.

The compound of formula IH can then be converted into the corresponding nitrogen protected derivative of formula XX, using the methods described above for synthesizing the compound of formula IV in Scheme 1. Nitration of the compound of formula XX using the
15 method described above for preparing the compound of formula IX in Scheme 6, yields the compound of formula XXI wherein the benzo ring is substituted with both a fluoro and nitro group or an alkoxy group and nitro group. The compound of formula XXI can be used to make a variety of compounds of the formula I wherein one of R² and R³ is fluoro, using methods that are well known to those of skill in the art, for example, by first converting the nitro group to an amino
20 group, converting the amino group to a variety of other substituents, as illustrated in Scheme 10, and then removing the nitrogen protecting group.

The compound of formula XXI acts as a regioisomeric functional equivalent of the compounds having formulas IIA, VIA and XXII, in that the fluorine atom of formula XXI reacts similarly to the nitro and Y groups of formula IIA, VIA, and XXII, and thus can be subjected to the
25 same series of reactions as those described above for the latter three compounds, providing an alternate means for preparing the products of such reactions. Similarly, the alkoxy group of formula XXI (R¹⁸=alkoxy) may be converted into a hydroxyl group before or after introduction of the nitro group, and then converted to isomeric products as described above. Also, the trifluoromethanesulfonate salt of such hydroxy derivative can act as a Y-group as described.

30 Preparation of compounds of formula I where R² = -O(C₁-C₆)alkyl, (C₁-C₆) alkyl or aryl wherein aryl is defined as above in the definition of formula I, and R³ is H or one of the other substituents described above in the definition of formula I, can be prepared as described above and illustrated in Scheme 8 by replacing one of the fluorine atoms of the compound of formula XV with -O-(C₁-C₆)alkyl, (C₁-C₆)alkyl or aryl, respectively

35 Scheme 9 illustrates methods of preparing compounds of the formula I wherein: (a) R¹ is hydrogen and R² is R⁷R⁸NO₂S-, (b) R¹ and R² are both chloro; and (c) R¹ is hydrogen and R² is R¹³C(=O)-. These compounds are referred to in Scheme 9, respectively, as compounds of formulas IJ, IK and IL.

5 Referring to Scheme 9, compounds of the formula IJ can be prepared by reacting the compound of formula IV with two or more equivalents of a halosulfonic acid, preferably chlorosulfonic acid, at a temperature from about 0°C to about room temperature. Reaction of the chlorosulfonic acid derivative so formed with an amine having the formula R^7R^8NH , wherein R^7 and R^8 are defined as above, followed by removal of the nitrogen protecting group,
10 yields the desired compound having formula IJ.

Compounds of the formula IK can be prepared by reacting the compound of formula IV with iodine trichloride in a chlorinated hydrocarbon solvent, followed by removal of the nitrogen protecting group. The reaction with iodine trichloride is typically carried out at a temperature from about 0°C to about room temperature, and is preferably carried out at about
15 room temperature. In a similar fashion, the analogous mono- or dibrominated or mono- or diiododinated compounds can be prepared by reacting the compound of IV with N-iodosuccinimide or N-bromosuccinimide in a trifluoromethanesulfonic acid solvent, followed by removal of the nitrogen protecting group as described above

Reaction of the compound of IV with an acid halide of the formula $R^{13}COCl$ or an acid
20 anhydride of the formula $(R^{13}CO)_2O$, with or without a reaction inert solvent such as a chlorinated hydrocarbon solvent, preferably methylene chloride, in the presence of Lewis acid such as aluminum chloride, at a temperature from about 0°C to about 100°C, followed by nitrogen deprotection, yields the compound of formula IL. The reaction with the acid halide or anhydride can be carried out using other known Lewis acids or other Friedel-Crafts acylations
25 methods that are known in the art.

The reactions described herein in which NO_2 , $-SO_2NR^7R^8$, $-COR^{13}$, I, Br or Cl are introduced on the compound of formula IV, as depicted in Scheme 9 and described above, can be performed on any analogous compound wherein R^2 is hydrogen, $(C_1-C_6)alkyl$, halo $(C_1-C_6)alkoxy$ or $-NHCONR^7R^8$, producing compounds of the formula I wherein R^2 and R^3 are
30 defined as in the definition of compounds of the formula I above.

Compounds that are identical to those of the formula IL, but which retain the nitrogen protecting group, can be converted into the corresponding O-acyl substituted compounds, i.e., those wherein the $-C(=O)R^{13}$ group of formula IL is replaced with a $-O-C(=O)R^{13}$ group, using Baeyer-Villiger processes well known to those skilled in the art. The resulting compounds can
35 be partially hydrolyzed, as described in Example 35, to yield the corresponding hydroxy substituted compounds, and then alkylated to form the corresponding alkoxy substituted compounds. Also, as described in Example 36, such O-acyl substituted compounds can be used to prepare variably substituted benzisoxazoles

5 Scheme 10 illustrates methods of making compounds of the formula I wherein: (a) R¹ is hydrogen and R² is chloro; (b) R¹ is hydrogen and R² is cyano, (c) R¹ is hydrogen and R² is amino; and (d) R¹ is hydrogen and R² is R¹³C(=O)N(H)-. These compounds are referred to in Scheme 10, respectively, as compounds of the formula IM, IN, IP and IQ.

10 Compounds of formula IM can be prepared from compounds of the formula IX' by generation of a diazonium salt with, for instance, an alkali metal nitrite and strong mineral acid (e.g., hydrochloric acid, sulfuric acid, hydrobromic acid) in water, followed by reaction with a copper halide salt, such as copper (I) chloride. Nitrogen deprotection by the methods described above yields the desired compound of formula IM. Alternative methods for the generation of diazonium salts, as known and practiced by those of skill in the art, can also be used. The foregoing reaction is generally carried out by temperatures ranging from about 0°C to about 60°C, preferably about 60°C for about 15 minutes to one hour.

20 Reaction of the diazonium salt, prepared as described above, with potassium iodide in an aqueous medium provides the analogous iodide derivative. This reaction is generally carried out at a temperature from about 0°C to about room temperature, preferably at about room temperature. The resulting compound, or its analogous N-tert-butylcarbonate protected form, can be used to prepare the corresponding cyano derivative by reaction with copper (I) cyanide and sodium cyanide in DMF, N,N-dimethylpropylurea (DMPU) or DMSO, preferably DMF, at a temperature from about 50°C to about 180°C, preferably about 150°C. Nitrogen deprotection as described above provides the desired compound of formula IM.

25 The above described iodide derivative can also be used to access a variety of other substituents such as aryl, acetylene and vinyl substituents, as well as the corresponding carbonyl esters and amides, by palladium and nickel catalyzed processes known to those of skill in the art, such as Heck, Suzuki and Stille couplings and Heck carbonylations.

30 Nitrogen deprotection of the compound of formula IX' provides the compound of the formula IP.

35 The compound of formula IX' can be reacted with a acyl group having the formula R¹³COCl or (R¹³CO)₂O using the methods described above, followed by nitrogen deprotection to provide compounds of the formula IQ. In a similar fashion, treatment of the protected amine with a compound having the formula R¹³SO₂X, when X is chloro or bromo, followed by nitrogen deprotection, provides the corresponding sulfonamide derivative.

 Other suitable amine protecting groups that can be used, alternatively, in the procedures described throughout this document include -COCF₃, -COCCl₃, -COOCH₂CCl₃, -COO(C₁-C₆)alkyl and -COOCH₂C₆H₅. These groups are stable under the conditions

5 described herein, and may be removed by methods described for each in Greene's "Protective Groups in Organic Chemistry", referred to above.

In each of the reactions discussed above, or illustrated in Schemes 1-10, above, pressure is not critical unless otherwise indicated. Pressures from about 0.5 atmospheres to about 5 atmospheres are generally acceptable, with ambient pressure, i.e., about 1 atmosphere, being preferred as a matter of convenience.

The compounds of the formula I and their pharmaceutically acceptable salts (hereafter "the active compounds") can be administered via either the oral, transdermal (e.g., through the use of a patch), intranasal, sublingual, rectal, parenteral or topical routes. Transdermal and oral administration are preferred. These compounds are, most desirably, administered in dosages ranging from about 0.25 mg up to about 1500 mg per day, preferably from about 0.25 to about 300 mg per day in single or divided doses, although variations will necessarily occur depending upon the weight and condition of the subject being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 0.01 mg to about 10 mg per kg of body weight per day is most desirably employed. Variations may nevertheless occur depending upon the weight and condition of the persons being treated and their individual responses to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval during which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effects, provided that such larger doses are first divided into several small doses for administration throughout the day.

The active compounds can be administered alone or in combination with pharmaceutically acceptable carriers or diluents by any of the several routes previously indicated. More particularly, the active compounds can be administered in a wide variety of different dosage forms, e.g., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, transdermal patches, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents. In addition, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the active compounds are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

5 For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc can be used for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar] as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration the active ingredient may be combined with various sweetening or flavoring agents, coloring matter and, if so desired, emulsifying and/or suspending agents, together with such diluents as water, ethanol, propylene glycol, glycerin and various combinations thereof.

For parenteral administration, a solution of an active compound in either sesame or peanut oil or in aqueous propylene glycol can be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8), if necessary, and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

It is also possible to administer the active compounds topically and this can be done by way of creams, a patch, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

Biological Assay

The effectiveness of the active compounds in suppressing nicotine binding to specific receptor sites is determined by the following procedure which is a modification of the methods of Lippiello, P. M. and Fernandes, K. G. (in The Binding of L-[³H]Nicotine To A Single Class of High-Affinity Sites in Rat Brain Membranes, Molecular Pharm, 29, 448-54, (1986)) and Anderson, D. J. and Arnenc, S. P. (in Nicotinic Receptor Binding of ³H-Cystisine, ³H-Nicotine and ³H-Methylcarbamylcholine In Rat Brain, European J. Pharm., 253, 261-67 (1994)).

5

Procedure

Male Sprague-Dawley rats (200-300 g) from Charles River were housed in groups in hanging stainless steel wire cages and were maintained on a 12 hour light/dark cycle (7 a.m.-7 p.m. light period). They received standard Purina Rat Chow and water *ad libitum*.

10 The rats were killed by decapitation. Brains were removed immediately following decapitation. Membranes were prepared from brain tissue according to the methods of Lippiello and Fernandez (Molec Pharmacol, 29, 448-454, (1986) with some modifications. Whole brains were removed, rinsed with ice-cold buffer, and homogenized at 0° in 10 volumes of buffer (w/v) using a Brinkmann Polytron™, setting 6, for 30 seconds. The buffer consisted of 50 mM Tris HCl at a pH of 7.5 at room temperature. The homogenate was sedimented by centrifugation (10 15 minutes; 50,000 x g; 0 to 4°C. The supernatant was poured off and the membranes were gently resuspended with the Polytron and centrifuged again (10 minutes; 50,000 x g; 0 to 4°C. After the second centrifugation, the membranes were resuspended in assay buffer at a concentration of 1 0g/100mL. The composition of the standard assay buffer was 50 mM Tris HCl, 120 mM NaCl, 5 mM KCl, 2 mM MgCl₂, 2 mM CaCl₂ and has a pH of 7.4 at room temperature.

20 Routine assays were performed in borosilicate glass test tubes. The assay mixture typically consisted of 0.9 mg of membrane protein in a final incubation volume of 1.0 mL. Three sets of tubes were prepared wherein the tubes in each set contained 50µL of vehicle, blank, or test compound solution, respectively. To each tube was added 200 µL of [³H]-nicotine in assay buffer followed by 750 µL of the membrane suspension. The final concentration of nicotine in 25 each tube was 0.9 nM. The final concentration of cytosine in the blank was 1 µM. The vehicle consisted of deionized water containing 30 µL of 1 N acetic acid per 50 mL of water. The test compounds and cytosine were dissolved in vehicle. Assays were initiated by vortexing after addition of the membrane suspension to the tube. The samples were incubated at 0 to 4° C in an iced shaking water bath. Incubations were terminated by rapid filtration under vacuum through 30 Whatman GF/B™ glass fiber filters using a Brandel™ multi-manifold tissue harvester. Following the initial filtration of the assay mixture, filters were washed two times with ice-cold assay buffer (5 mL each). The filters were then placed in counting vials and mixed vigorously with 20 mL of Ready Safe™ (Beckman) before quantification of radioactivity. Samples were counted in a LKB Wallach Rackbeta™ liquid scintillation counter at 40-50% efficiency. All determinations were in 35 triplicate.

5

Calculations

Specific binding (C) to the membrane is the difference between total binding in the samples containing vehicle only and membrane (A) and non-specific binding in the samples containing the membrane and cytosine (B), i.e.,

$$\text{Specific binding} = (C) = (A) - (B).$$

10

Specific binding in the presence of the test compound (E) is the difference between the total binding in the presence of the test compound (D) and non-specific binding (B), i.e., $(E) = (D) - (B)$.

$$\% \text{ Inhibition} = (1 - ((E)/(C))) \text{ times } 100.$$

15 The compounds of the invention that were tested in the above assay exhibited IC_{50} values of less than $10 \mu M$.

The following experimental examples illustrate, but do not limit the scope of, this invention

EXAMPLE 1

10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE

20

A) 1,4-Dihydro-1,4-methano-naphthalene

(Based wholly or in part on a) Wittig, G.; Knauss, E. *Chem Ber.* **1958**, 91, 895. b) Muir, D. J., Stothers, J. B. *Can. J. Chem.* **1993**, 71, 1290.)

25 Magnesium turnings (36.5 g, 1.5 M) were stirred in anhydrous THF (250 mL) in a dried 2 L 3 neck round bottom flask equipped with a 250 mL non-equalizing addition funnel with a nitrogen (N_2) flow adapter, mechanical stirrer and efficient condenser equipped with a N_2 flow adapter. The flask was stirred and warmed to reflux by a removable heating mantle. 2-Fluorobromobenzene (2g) was added followed by 1 mL of 3N ethylmagnesium bromide (EtMgBr in THF). The addition funnel was charged with a mixture of cyclopentadiene (94.4 g, 1.43 M. Prepared by the method described in: *Org. Syn. Col. Vol. V*, 414-418) and 30 bromofluorobenzene (250 g, 1.43 M) which was maintained at $0^\circ C$ in a separate flask by an ice bath, and transferred to the addition funnel via cannula. Small portions (~1 mL) of the intimate mixture were introduced to assist initiation (~4x). After ~15 minutes, the reaction initiated (exotherm, and vapor condensation), the heating mantle was removed and the contents of the addition funnel was added dropwise at such rate as to maintain reflux (1.5 35 hours). The heating mantle was re-applied and a reflux maintained for 1.5 hours. (TLC 100% hexanes R_f 0.67)

The reaction was cooled to room temperature and quenched with H_2O (500 mL) and carefully with 1N HCl (200 mL, produces H_2 evolution from unconsumed Mg). To this ~50 mL

- 5 concentrated HCl was added to dissolve solids. Total addition/quench time ~1 hour
Saturated aqueous sodium chloride (NaCl) solution (300mL) was added and product hexanes
extracted until no potassium permanganate (KMnO₄) active product is removed. (4 x ~250
mL). The combined organic layer was washed with saturated NaHCO₃ solution (250 mL),
sodium bicarbonate Na₂SO₄ dried and concentrated to an oil (~200 g). The product was
10 distilled at 78-83 °C @15mm (131 g, 64%). (An alternative workup is described on p.419
Fieser and Fieser, Vol. I, Reagents for Organic Synthesis, Wiley, NY., NY.; 1967).

B) 1,2,3,4-Tetrahydro-1,4-methano-naphthalene-2,3-diol

- (Except for the workup method and the quantity of OsO₄ used, based on
15 VanRheenen, V ; Cha, D.Y.; Hartley, W M. *Org. Syn* **1988**, 6, 342.)

In a 2 L 3 neck round bottom flask equipped with a N₂ flow adapter, mechanical stirrer
was placed 1,4-dihydro-1,4-methano-naphthalene (79.5 g, 560 mmol) stirred in acetone (800
mL) and H₂O (100 mL) and N-methyl morpholine N-oxide (67.5 g, 576 mmol). To this was
added osmium tetroxide (OsO₄) (15 mL of a 15mol% t-BuOH solution, 1.48 mmol, 0.26mol%)
20 and the mixture was stirred vigorously. After 60 hours, the reaction was filtered, and the white
product rinsed with acetone and air dried (60.9 g). The mother liquor was concentrated to an
oily solid; acetone trituration, filtration and acetone rinse provided (27.4 g, total 88.3 g, 89%)
(TLC 50% EtOAc/hexanes R_f ~0.5). mp 176-177.5 °C.

- 25 C) 10-Benzyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

(Based on Abdel-Magid, A. F., Carson, K. G., Harris, B. D.; Maryanoff, C. A.; Shah, R.
D. *J. Org. Chem.* **1996**, 61, 3849, and Mazzocchi, P. H.; Stahly, B. C. *J. Med. Chem.* **1979**,
22, 455.)

- 1,2,3,4-Tetrahydro-1,4-methano-naphthalene-2,3-diol (40 g, 227.3 mmol) was stirred
30 in H₂O (1050 mL) and 1,2-dichloroethane (DCE) (420 mL) in a 2 L round bottom flask under
nitrogen with cool water bath (~10 °C). To this sodium periodate (NaIO₄) (51 g, 239 mmol)
and triethylbenzyl ammonium chloride (Et₃BnNCl) (50 mg) were added. The resulting mixture
was stirred for 1 hour (slight initial exotherm), then the layers were separated and the
aqueous layer was extracted with DCE (200 mL). The organic layer was washed with H₂O (4
35 x 200 mL, or until no reaction to starch iodide is observed in the aqueous wash) then dried
through a cotton plug. To this was added benzyl amine (25.5 g, 238.6 mmol) and the mixture
was stirred for 2 minutes then immediately transferred into the sodium triacetoxyborohydride
NaBH(OAc)₃/DCE (see below) over 10 minutes

- 5 In a separate 2 L round bottom flask under nitrogen was magnetically stirred NaHB(OAc)₃ (154 g, 0.727 mmol) in DCE (800 mL) at 0 °C (ice bath). To this was added the above mixture over 10 minutes, without delay after the dialdehyde and amine were mixed. The resulting orange mixture was allowed to warm to room temperature and stirred for 30-60 minutes.
- 10 The reaction was quenched by addition of saturated sodium carbonate (Na₂CO₃) solution (~300 mL) carefully at first and the mixture was stirred for 1 hour (pH 9). The layers were separated and the aqueous layer was extracted with CH₂Cl₂ (2 x 300 mL). The organic layer was washed with saturated aqueous NaCl solution (200 mL), dried through a cotton plug, then evaporated to a red oil. This was dissolved in a minimum of Et₂O and filtered
- 15 through a Silica pad (3 x 4 inch) eluting with 15% ethyl acetate (EtOAc)/hexanes +1% of 37% aqueous ammonium hydroxide (NH₄OH) solution to remove baseline red color. Concentration affords a light yellow oil (48.5 g, 194.8 mmol, 85.7%). (TLC 10% EtOAc/hexanes R_f 0.75). ¹H NMR (400 MHz, CDCl₃) δ 7.16 (m, 7H), 6.89 (m, 2H), 3.48 (m, 2H), 3.08 (m, 2H), 2.80 (d, J=9.5 Hz, 2H), 2.42 (d, J=9.5 Hz, 2H), 2.27 (m, 1H), 1.67 (d, J=10.0 Hz, 1H). APCI MS m/e
- 20 250.3 [(M + 1)⁺].

D) 10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (For an alternative synthesis, see; Mazzocchi, P. H.; Stahly, B. C. *J. Med. Chem.* **1979**, 22, 455.)

- 10-Benzyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (70.65 g, 284 mmol) was
- 25 stirred in EtOAc (250 mL) and treated with 3N HCl EtOAc (1.03 eq.) slowly with cooling (ice bath). The resulting precipitate was filtered and rinsed with EtOAc. The solids were dissolved in MeOH (250 mL) in a Parr bottle. To this was added Pd(OH)₂ (7 g of 20%wt/C) and the mixture was shaken under 50-40 psi of H₂ for 24 hours or until done by TLC. The reaction was filtered through a Celite pad and concentrated to an oily solid. This was azeotroped with
- 30 methanol (MeOH) (3x) then triturated with acetone, treated with ethyl ether (Et₂O) to precipitate product and filtered. Concentration of the mother liquors and a second treatment provided an off white solid (48.95 g, 251 mmol, 88%). (TLC 10% MeOH/CH₂Cl₂ (NH₃) R_f 0.2). ¹H NMR (400 MHz, CDCl₃) δ 7.18 (m, 4H), 2.97 (m, 4H), 2.68 (d, J=12.5 Hz, 2H), 2.41 (m, 1H), 1.95 (d, J=11.0 Hz, 1H). APCI MS m/e 160.2 [(M + 1)⁺].

5

EXAMPLE 2

4-FLUORO-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7).3,5-TRIENE
HYDROCHLORIDE

A) 6-Fluoro-1,4-dihydro-1,4-methano-naphthalene

(Eisch, J. J.; Burlinson, N. E. *J. Amer. Chem. Soc.* **1976**, *98*, 753-761. Paquette, L. A.;

10 Cottrell, D. M.; Snow, R. A. *J. Amer. Chem. Soc.* **1977**, *99*, 3723-3733.)

Magnesium turnings (0.66 g, 27.2 mmol) were stirred in anhydrous THF (10 mL) in a flame dried 75 mL 3 neck round bottom flask equipped with a non-equalizing addition funnel with a N₂ flow adapter, magnetic stirrer and efficient condenser equipped with a N₂ flow adapter. The flask was stirred and warmed to reflux by a removable heating mantle. 2,5-
15 Difluorobromobenzene (0.1 g) was added followed by of 3N EtMgBr in THF (0.1 mL). The addition funnel was charged with an intimate mixture of cyclopentadiene (1.71 g, 25.9 mmol) and 2,5-difluorobromobenzene (5.0 g, 25.9 mmol). Small portions (~0.2 mL) of the intimate mixture were introduced to assist initiation (~4x). After ~15 minutes, the reaction initiated (exotherm, and vapor condensation) and heating was maintained as necessary during the
20 addition of the contents of the addition funnel. The reaction was then maintained at reflux for 1 hour

The reaction was cooled to room temperature and quenched with H₂O (20 mL) followed by aqueous 1N HCl solution (20 mL) to dissolve the solids. Saturated aqueous NaCl solution (30 mL) was added and product was extracted with hexanes (4 x 25mL). The
25 combined organic layer was washed with saturated aqueous NaHCO₃ solution (25 mL), dried (Na₂SO₄), filtered through a Silica plug with hexanes rinse and concentrated to an oil. Chromatography on Silica gel eluting with hexanes provided an oil (780 mg, 19%). (TLC hexanes R_f 0.38). ¹H NMR (400 MHz, CDCl₃) δ 7.10 (m, 1H), 6.97 (d, J=8.0 Hz, 1H), 6.80 (br s, 1H), 6.78 (br s, 1H), 6.59 (m, 1H), 3.87 (br s, 2H), 2.32 (d, J=7.0 Hz, 1H), 2.25 (d, J=7.0 Hz,
30 1H)

B) 6-Fluoro-1,2,3,4-tetrahydro-1,4-methano-naphthalene-2,3-diol

6-Fluoro-1,4-dihydro-1,4-methano-naphthalene (680 mg, 4.22 mmol) and N-methyl morpholine N-oxide (599 mg, 4.43 mmol) were stirred in acetone (50 mL) and H₂O (5 mL). To
35 this was added a solution of OsO₄ (0.2 mL, 2.5%wt. solution in t-BuOH, 0.02 mmol). After 72 hours, florisil (5 g) and saturated aqueous NaHSO₃ solution (3 mL) were added and stirred for 1 hour. The florisil was filtered and the filtrate concentrated to produce a crystalline product which was triturated with acetone and filtered (524 mg, 64%). ¹H NMR (400 MHz, CDCl₃) δ

09202010 09202010

5 7.10 (dd, J=8.0,5.0 Hz, 1H), 6.90 (dd, J=8.0,2.3 Hz, 1H), 6.75 (ddd, J=8.0,8.0,2.3 Hz, 1H),
3.79 (s, 2H), 3.18 (d, J=1.5 Hz, 2H), 2.22 (d, J=10.0 Hz, 1H), 1.92 (dd, J=10.0,1.5 Hz, 1H).
GCMS *m/e* 194 (*M*⁺)

C) 10-Benzyl-4-fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

10 6-Fluoro-1,2,3,4-tetrahydro-1,4-methano-naphthalene-2,3-diol (524 mg, 2.68 mmol)
and Et₃NBnCl (10 mg) were vigorously stirred in dichloroethane (15 mL) and H₂O (45 mL)
then treated with sodium periodate (0.603 mg, 2.82 mmol). After 1.5 hours, the layers were
separated and the aqueous layer extracted with DCE (2 x 20 mL). The combined organic
layer was washed with H₂O (4 x 20 mL) until no reaction to starch iodide paper was observed,
15 then with saturated aqueous NaCl solution (20 mL). The organic layer was dried through a
cotton plug and treated with benzyl amine (0.308 mL, 2.82 mmol) and stirred for 2 minutes
then transferred to an addition funnel. This solution was added over ~10 minutes to a
vigorously stirred cooled (0 °C) mixture of NaHB(OAc)₃ (1.82 g, 8.58 mmol) in DCE (50 mL).
After addition was complete, the mixture was stirred without cooling for 2 hours. The mixture
20 was quenched with saturated aqueous Na₂CO₃ solution (100 mL) and stirred for 1 hour, then
the layers were separated and the aqueous layer was extracted with CH₂Cl₂ (3 x 30 mL). The
combined organic layer was washed with saturated aqueous NaCl solution (50 mL), dried
through a cotton plug and concentrated. Chromatography on Silica gel provided an oil (520
mg, 80%). (TLC 2% acetone/CH₂Cl₂ R_f 0.40). ¹H NMR (400 MHz, CDCl₃) δ 7.18 (m, 1H), 6.88
25 (m, 2H), 3.48 (s, 2H), 3.06 (m, 2H), 2.78 (m, 2H), 2.41 (m, 2H), 2.27 (m, 1H), 1.69 (d, J=10.5
Hz, 1H).

D) 4-Fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

10-Benzyl-4-fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (390 mg, 1.461
30 mmol), ammonium formate (3.04 g, 48.2 mmol) and 10%Pd(OH)₂/C (30 mg) were combined in
MeOH (50 mL) and brought to reflux under N₂ for 1.5 hours. Ammonium formate (1.0 g) was
added and reflux continued for 0.5 hour. The reaction mixture was filtered through a Celite
pad which was rinsed with MeOH. The filtrate was concentrated. The residues were treated
with saturated aqueous Na₂CO₃ solution (30 mL) and product extracted with methylene
35 chloride (CH₂Cl₂) (3 x 25 mL). The organic layer was washed with saturated aqueous NaCl
solution (50 mL), dried through a cotton plug and concentrated. The residue was treated with
2N HCl MeOH (5 mL) and concentrated then taken up in minimum of MeOH and saturated
with Et₂O. After stirring 18h, the white crystals were collected by filtration (86 mg, 28%) (TLC

00402010 09369
00000000 00000000

- 5 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.27) (data for free base) ¹H NMR (400 MHz, CDCl₃) δ 7.06 (m, 1H), 6.83 (m, 2H), 2.89 (m, 4H), 2.61 (dd J=12.0 Hz, 2H), 2.37 (m, 1H), 1.87 (d, J=11.5 Hz, 1H). APCI MS *m/e* 178.2 [(M + 1)⁺]. (HCl salt) mp 260-262 °C.

EXAMPLE 3

10 4-METHYL-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE
HYDROCHLORIDE

- The title compound was prepared by the methods described in Example 1 and 2 starting with 2-fluoro-5-methylbromobenzene. (data for free base) ¹H NMR (400 MHz, CDCl₃) δ 7.04 (d, J=7.5 Hz, 1H), 6.99 (s, 1H), 6.98 (d, J=7.5 Hz, 1H), 2.98-2.90 (m, 4H), 2.63 (m, 2H),
15 2.35 (m, 1H), 2.32 (s, 3H), 1.87 (d, J=11.5 Hz, 1H) APCI MS *m/e* 174.2 [(M + 1)⁺]. (HCl salt) mp 254-255 °C. Anal. Calcd. for C₁₂H₁₂F₃N HCl.1/3H₂O C, 53.44; H, 5.11; N, 5.19. Found C, 53.73; H, 4.82; N, 5.15.

EXAMPLE 4

- 4-TRIFLUOROMETHYL-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE
20 HYDROCHLORIDE (See Grunewald, G. L.; Paradkar, V. M.; Pazhenchevsky, B.; Pleiss, M. A.; Sall, D. J.; Seibel, W. L.; Reitz, T. J. *J. Org. Chem.* **1983**, *48*, 2321-2327 Grunewald, G. L.; Markovich, K. M.; Sall, D. J. *J. Med. Chem.* **1987**, *30*, 2191-2208.)

- The title compound was prepared by the methods described in Example 1 and 2 starting with 2-fluoro-5-trifluoromethylbromobenzene ¹H NMR (400 MHz, CD₃OD) δ 7.71 (s, 1H), 7.64 (d, J=8.0 Hz, 1H), 7.57 (d, J=8.0 Hz, 1H) 3.46 (m, 4H), 3.21 (d, J=12.5 Hz, 2H),
25 2.41 (m, 1H), 2.16 (d, J=11.5 Hz, 1H) APCI MS *m/e* 228.2 [(M + 1)⁺] (HCl salt) mp 244-246 °C. Anal. Calcd. for C₁₂H₁₂F₃N HCl.1/3H₂O C, 53.44; H, 5.11; N, 5.19. Found C, 53.77; H, 4.82; N, 5.18

30 EXAMPLE 5

- 3-TRIFLUOROMETHYL-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE
HYDROCHLORIDE (Grunewald, G. L.; Markovich, K. M.; Sall, D. J. *J. Med. Chem.* **1987**, *30*, 2191-2208)

- The title compound was prepared by the methods described in Example 1 and 2 starting with 2-fluoro-6-trifluoromethylbromobenzene ¹H NMR (400 MHz, CD₃OD) δ 7.65 (s, 2H), 7.52 (m, 1H), 3.65 (br s, 1H), 3.49-3.43 (m, 3H), 3.20 (m, 2H), 2.42 (m, 1H), 2.18 (d, J=11.5 Hz, 1H) APCI MS *m/e* 228.2 [(M + 1)⁺] (HCl salt) mp 275-277 °C.
- 35

5

EXAMPLE 6

3-FLUORO-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE

HYDROCHLORIDE

A) 2,6-Difluoriodobenzene (Roe, A. M.; Burton, R. A.; Willey, G. L.; Baines, M. W.; Rasmussen, A. C. *J. Med. Chem.* **1968**, 11, 814-819. Tamborski, C.; Soloski, E. *J. Org. Chem.* **1966**, 31, 746-749. Grunewald, G. L.; Arrington, H. S.; Bartlett, W. J.; Reitz, T. J.; Sall, D. J. *J. Med. Chem.* **1986**, 29, 1972-1982.) 1,3-Difluorobenzene (57.05 g, 0.5 M) in THF (75 mL) was added to a -78 °C stirred solution of n-butyllithium (n-BuLi) (200 mL, 2.5 M/hexanes, 0.5 M) and THF (500 mL) under N₂. By controlling the addition rate the internal temperature was maintained below -70 °C. The total addition time was ~1/2 hour. The resulting slurry was stirred an additional 1/2 hour, then the dispersion was treated with a solution of iodine (126.9 g, 0.5 M) in THF (300 mL) at a rate that maintained an internal temperature below -70 °C. After complete addition the mixture was allowed to warm to room temperature, and was treated with H₂O (100 mL) and 10% aqueous Na₂S₂O₃ solution (100 mL) and stirred. The layers were separated and the aqueous layer extracted with hexanes (2 x 250 mL). The combined organic layer was washed with 10% aqueous Na₂S₂O₃ solution (100 mL), H₂O (100 mL), saturated aqueous NaCl solution (100 mL), dried (Na₂SO₄) filtered and concentrated to give a yellow oil (106.5 g). Distillation at ~1-5 mm at ~80 °C provided a light yellow oil (89.5 g, 75%). ¹H NMR (400 MHz, CDCl₃) δ 7.30 (m, 1H), 6.87 (m, 2H) GCMS *m/e* 240 (M⁺).

25

B) 5-Fluoro-1,4-dihydro-1,4-methano-naphthalene

A solution of 2,6-difluoriodobenzene (5.0 g, 20.8 mmol) and cyclopentadiene (2.07 g, 31.3 mmol) was stirred at 0 °C in P. ether (70 mL, 40-60 °C) under N₂ and treated with n-BuLi (8.74 mL, 2.5M in hexanes, 21.8 mmol) dropwise over 10 minutes. The reaction was quenched after 15 minutes by addition of aqueous 1N HCl solution and the product was extracted with hexanes (3 x 50 mL). The combined organic layer was washed with H₂O (50 mL), saturated aqueous NaCl solution (50 mL), dried (MgSO₄), filtered and evaporated. Chromatography on Silica gel provided product as an oil (1.5 g, 45%) (TLC hexanes R_f 0.55). ¹H NMR (400 MHz, CDCl₃) δ 7.08 (ddd, J=7.0,1.0,0.8 Hz, 1H), 6.96 (ddd, J=8.5,8.3,7.0 Hz, 1H), 6.86 (br s, 2H), 6.72 (ddd, J=8.5,8.3,0.8 Hz, 1H), 4.25 (br s, 1H), 3.98 (br s, 1H), 2.36 (ddd, J=7.2,1.7,1.7 Hz, 1H), 2.30 (ddd, J=7.2,1.7,1.5 Hz, 1H). GCMS *m/e* 160 (M⁺).

35

00402010 0920460

5 C) 3-Fluoro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

The title compound was prepared by the methods described in Example 2B,C,D starting with 5-fluoro-1,4-dihydro-1,4-methano-naphthalene. ¹H NMR (400 MHz, CD₃OD) δ 7.36 (ddd J=8.3,7.3,5.0 Hz, 1H), 7.21 (d, J=7.3 Hz, 1H), 7.07 (t, J=8.3 Hz, 1H), 3.62 (br s, 1H), 3.42-3.30 (m, 3H), 3.21 (m, 2H), 2.38 (m, 1H), 2.12 (d, J=11.5 Hz, 1H). APCI MS *m/e* 178.4 [(M + 1)⁺]. mp 269-271 °C.

EXAMPLE 7

4-NITRO-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE
HYDROCHLORIDE

A) 1-(10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone

15 10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride salt (12.4 g, 63.9 mmol) was stirred in CH₂Cl₂ (200 mL). This was cooled (ice bath) and treated with pyridine (12.65 g 160 mmol) followed by trifluoroacetic anhydride (TFAA) (16.8 g, 11.3 mL, 80 mmol) from an addition funnel over 10 minutes. After ~3 hours, the solution was poured into 0.5N aqueous HCl (200 mL) and the layers separated. The aqueous layer was extracted with
20 CH₂Cl₂ (3 x 50 mL) and the combined organic layer was washed with 0.5N aqueous HCl (50 mL), H₂O (2 x 50 mL) and saturated aqueous NaHCO₃ solution (50 mL). This solution was dried through a cotton plug, then diluted with ~3% EtOAc and filtered through a 2 inch Silica pad eluted with ~3% EtOAc/CH₂Cl₂. Concentration afforded a clear oil which crystallized to give white needles (15.35 g, 60.2 mmol, 94%). (TLC 30% EtOAc/hexanes R_f 0.53). ¹H NMR
25 (400 MHz CDCl₃) δ 7.18 (m, 4H), 4.29 (br d, J=12.6 Hz, 1H), 3.84 (br d, J=12.6 Hz, 1H), 3.51 (dd, J=12.6,1.5 Hz, 1H), 3.21 (br s, 1H), 3.10 (br s, 1H), 3.10 (br d, J=12.6 Hz, 1H), 2.37 (m, 1H), 1.92 (d, J=10.8 Hz, 1H). GCMS *m/e* 255 (M⁺). mp 67-68 °C.

B) 1-(4-Nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
30 ethanone (Based on the method described by Coon, C. L., Blucher, W G, Hill, M. E. *J Org Chem.* **1973**, 25, 4243)

To a solution of trifluoromethanesulfonic acid (2.4 mL, 13.7 mmol) in CH₂Cl₂ (10 mL) stirred at 0 °C was slowly added nitric acid (0.58 mL, 27.4 mmol) generating a white precipitate. After 10 minutes the resulting mixture was cooled to -78 °C and treated with 1-
35 (10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (3.5 g, 13.7 mmol) in CH₂Cl₂ (15 mL) dropwise from an addition funnel over 5 minutes. The reaction was stirred at -78 °C for 30 minutes then warmed to 0 °C for 1 hour. The reaction mixture was poured into a vigorously stirred ice (100 g). The layers were separated and the aqueous layer

5 extracted with CH₂Cl₂ (3 x 30 ml). The organic layer was combined and washed with H₂O (3 x 30 ml). The combined organic layer was washed with saturated aqueous NaHCO₃ solution (20 mL) and H₂O (20 mL) then dried through a cotton plug and concentrated to give an orange oil that solidified on standing (4.2 g). Chromatography yielded pure product as a crystalline solid (3.2 g, 78%). (TLC 30% EtOAc/hexanes R_f 0.23). ¹H NMR (400 MHz, CDCl₃) δ 8.12 (br d, J=8.0 Hz, 1H), 8.08 (br s, 1H), 7.37 (br d, J=8.0 Hz, 1H), 4.38 (br d, J=12.6 Hz, 1H), 3.94 (br d, J=12.6 Hz, 1H), 3.59 (br d, J=12.6 Hz, 1H), 3.43-3.35 (m, 2H), 3.18 (br d, J=12.6 Hz, 1H), 2.48 (m, 1H), 2.07 (d, J=10.8 Hz, 1H). GCMS *m/e* 300 (M⁺).

C) 4-Nitro-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

15 1-(4-Nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (182 mg, 0.61 mmol) was stirred with Na₂CO₃ (160 mg, 1.21 mmol) in MeOH (3 mL) and H₂O (1 mL) at 70 °C for 18 hours. The mixture was concentrated, water was added and the product was extracted with CH₂Cl₂. The organic layer was extracted with 1N aqueous HCl (3 x 20 mL) and the acidic layer washed with CH₂Cl₂ (2 x 20 mL). The aqueous layer was
20 basified to pH ~10 with Na₂CO₃(s) and product was extracted with CH₂Cl₂ (3 x 30 mL). The organic layer was dried through a cotton plug and concentrated to an oil. This was dissolved in MeOH and treated with 1N HCl MeOH, concentrated to solids which were recrystallized from MeOH/Et₂O to afford product as a white solid (73 mg, 50%). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.38). ¹H NMR (400 MHz, DMSO-d₆) δ 8.21 (s, 1H), 8.18 (dd, J=8.0,2.0 Hz, 1H), 7.59 (d, J=8.0 Hz, 1H), 3.43 (br s, 2H), 3.28 (m, 2H), 3.07 (dd, J= 13.0,13.0 Hz, 2H), 2.24 (m, 1H),
25 2.08 (d, J=11.5 Hz, 1H) APCI MS *m/e* 205.1 [(M + 1)⁺] mp 265-270 °C

EXAMPLE 8

4-AMINO-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7).3.5-TRIENE

30 HYDROCHLORIDE

4-Nitro-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (500 mg, 2.08 mmol) was stirred in 1,4-dioxane (40 mL) and treated with saturated aqueous Na₂CO₃ solution (15 mL). To this was added di-*t*-butyldicarbonate (1.8 g, 8.31 mmol). After stirring 18 hours the reaction was treated with H₂O (50 mL), extracted with CH₂Cl₂ (4 x 30 mL), dried through a
35 cotton plug and concentrated to provide an oil (500 mg, 91%)

This oil (500 mg, 1.64 mmol) was dissolved in MeOH (30 mL), treated with 10%Pd/C (~50 mg) and hydrogenated under a H₂ atmosphere (45 psi) for 1 hour. The mixture was filtered through a Celite pad and concentrated to a clear oil (397 mg, 88%)

5 This oil (50 mg, 0.18 mmol) was stirred in 3N HCl EtOAc (3 mL) for 2 hours then concentrated to a white solid (25 mg, 56%). ¹H NMR (400 MHz, DMSO-d₆) δ 7.38-7.10 (3H), 3.60 (br s, 2H), 3.25 (m, 2H), 2.98 (m, 2H), 2.18 (m, 1H), 1.98 (d, J=11.5 Hz, 1H). APCI MS m/e 175.1 [(M + 1)⁺] mp 189-192 °C.

10

EXAMPLE 9

N¹-[10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YL]ACETAMIDE HYDROCHLORIDE

A) 1-(4-Amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone

15 Hydrogenation of 1-(4-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (2.0 g, 6.66 mmol) under a H₂ atmosphere (40 psi) and 10%Pd/C (200 mg) in MeOH over 1.5 hours, filtration through Celite and concentration affords a yellow oil (1.7 g). (TLC 50% EtOAc/hexanes R_f 0.27). ¹H NMR (400 MHz, CDCl₃) δ 6.99 (m, 1H), 6.64 (br s, 1H), 6.57 (m, 1H), 4.25 (m, 1H), 3.82 (m, 1H), 3.50 (m, 1H), 3.17-3.07 (m, 3H), 2.35 (m, 20 1H), 1.90 (d, J=10.8 Hz, 1H). GCMS m/e 270 (M⁺).

B) N-(10-Trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-acetamide

25 1-(4-Amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (850 mg, 3.14 mmol) was stirred in CH₂Cl₂ (5 mL) and treated with triethyl amine (0.53 mL, 3.76 mmol) and acetyl chloride (0.23 mL, 3.2 mmol) then stirred 18 hours. Standard NaHCO₃ workup yielded an oil which was chromatographed to provide a clear oil (850 mg, 87%). (50% EtOAc/hexanes R_f 0.28).

30 C) N¹-[10-Azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl]acetamide hydrochloride

N-(10-Trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-acetamide (100 mg, 0.32 mmol) was stirred with Na₂CO₃ (70 mg, 0.64 mmol) in MeOH (10 mL) and H₂O (2 mL) at 70 °C for 18 hours. The mixture was concentrated, water was added and the product was extracted with EtOAc. The organic layer was extracted with 1N aqueous HCl (3 x 35 20 mL) and the acidic layer washed with EtOAc (2 x 20 mL). The aqueous layer was basified to pH ~10 with Na₂CO₃ (s) and product was extracted with EtOAc (3 x 20 mL). The organic layer was dried (sodium sulfate (Na₂SO₄)) and concentrated to an oil. This material was dissolved in MeOH and treated with 3N HCl EtOAc (3 mL), concentrated and recrystallized

5 from MeOH/Et₂O to provide a solid (40 mg, 50%). ¹H NMR (400 MHz, DMSO-d₆) δ 9.98 (s, 1H), 9.02 (br m, NH), 7.65 (s, 1H), 7.55 (br s, NH), 7.38 (d, J=8.0 Hz, 1H), 7.20 (d, J=8.0 Hz, 1H), 3.33 (m, 4H), 2.96 (m, 2H), 2.13 (m, 1H), 2.00 (s, 3H), 1.96 (d, J=10.5 Hz, 1H). APCI MS m/e 217.2 [(M + 1)⁺]. mp 225-230 °C.

10

EXAMPLE 10

6-METHYL-5-THIA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,6,8-TETRAENE HYDROCHLORIDE

A) N-(10-Trifluorothioacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-thioacetamide

15 N-(10-Trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-acetamide (850 mg, 2.72 mmol) and 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's reagent) (1.1 g, 2.72 mmol) were combined in toluene (10 mL) and brought to reflux for 1.5 hours. After cooling the reaction was worked up with EtOAc/saturated aqueous NaHCO₃ solution. The organic layer was dried (Na₂SO₄), filtered, concentrated and
20 chromatographed on Silica gel to produce product (410 mg, 44%) (50% EtOAc/hexanes R_f 0.38)

B) 6-Methyl-5-thia-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,6,8-tetraene hydrochloride

25 The above oil, 2,2,2-trifluoro-N-(10-trifluorothioacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-thioacetamide, (360 mg, 1.05 mmol) was dissolved in MeOH (10 mL) and 1N NaOH (5 mL) and added to potassium ferricyanide (K₃Fe(CN)₆) (1.72 g, 5.23 mmol) in H₂O (10 mL). This mixture was warmed to 60 °C for 1.5 hours, cooled, concentrated and worked up with EtOAc/H₂O. This material was stirred in
30 dioxane (20 mL) and treated with H₂O (50 mL) and Na₂CO₃ to achieve pH 10. To this was added di-t-butylidicarbonate (436 mg, 2.0 mmol) and the mixture was stirred for 18 hours. The reaction was concentrated, treated with H₂O and extracted with CH₂Cl₂. The product was chromatographed (Silica 30% EtOAc/hexanes R_f 0.41) to yield an oil (100 mg).

The above product was treated with 3N HCl/EtOAc (3 mL) and warmed to reflux for
35 ~15 minutes then concentrated to a solid which was azeotroped with CH₂Cl₂ (2x). These solids were dissolved in a minimum amount of MeOH then saturated with Et₂O and stirred. The resulting white crystalline powder was collected by filtration (40 mg, 14%).

- 5 ¹H NMR (400 MHz, DMSO-d₆) δ 9.46 (s, NH), 7.65 (s, 1H), 7.82 (s, 1H), 7.65 (br m, NH), 3.36 (m, 2H), 3.24 (m, 2H), 3.02 (m, 2H), 2.76 (s, 3H), 2.23 (m, 1H), 2.06 (d, J=10.8 Hz, 1H). APCI MS *m/e* 231.1 [(M + 1)⁺] mp 183-184 °C.

EXAMPLE 11

10 4,5-DINITRO-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE

A) 1-(4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
ethanone (Based on the method described in Coon, C. L.; Blucher, W. G.; Hill, M. E. *J. Org. Chem.* **1973**, 25, 4243. For an additional related example of dinitration see: Tanida, H.; Ishitobi, H.; Irie, T.; Tsushima, T. *J. Am. Chem. Soc.* **1969**, 91, 4512.)

- 15 To a solution of trifluoromethanesulfonic acid (79.8 ml, 902.1 mmol) in CH₂Cl₂ (550 ml) stirred at 0 °C was slowly added nitric acid (19.1 ml, 450.9 mmol) generating a white precipitate. After 10 minutes, 1-(10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (50 g, 196 mmol) in CH₂Cl₂ (300 ml) was added dropwise from an addition funnel over 30 minutes. The reaction was stirred at 0 °C for 2.5 hours and then stirred at
 20 room temperature for 24 hours. The reaction mixture was poured into a vigorously stirred mixture of H₂O (500 ml) and ice (400 g). The layers were separated and the aqueous layer back extracted with CH₂Cl₂ (3 x 300 ml). The organic layer was combined and washed with H₂O (3 x 300 ml). The combined aqueous layers were re-extracted with CH₂Cl₂ (2 x 100 ml). The organic layer was combined and washed with saturated aqueous NaHCO₃ solution (200
 25 mL) and H₂O (200 mL) then dried through a cotton plug and concentrated to solids. Trituration with EtOAc/hexanes produced off white solids which were filtered and dried (52 g, 151 mmol, 77%). The mother liquor was chromatographed to give an additional 4.0 g for a total of 56.0 g (82.8%). (TLC 50% EtOAc/hexanes R_f 0.29) ¹H NMR (400 MHz, CDCl₃) δ 7.77 (s, 1H), 7.75 (s, 1H), 4.39 (br d, J=13.0 Hz, 1H), 3.98 (br d, J=13.0 Hz, 1H), 3.65 (d, J=13.0
 30 Hz, 1H), 3.49 (br s, 1H), 3.44 (br s, 1H), 3.24 (br d, J=12.6 Hz, 1H), 2.53 (m, 1H), 2.14 (d, J=11.5 Hz, 1H). GCMS *m/e* 345 (M⁺)

B) 4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

- 1-(4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
 35 ethanone (3.7 g, 10.7 mmol) and Na₂CO₃ (2.3 g, 21.4 mmol) were combined in MeOH (50 mL) and H₂O (20 mL) then warmed to reflux for 18 hours. The reaction was cooled, concentrated, treated with H₂O and extracted with CH₂Cl₂ (3 x 50 mL) then dried through a cotton plug. After concentration, the residue was chromatographed to provide brown solids. (1.9 g, 71%)

05402010 092839 152250 01020450

- 5 (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.36). ¹H NMR (400 MHz, CDCl₃) δ 7.69 (s, 2H), 3.17 (br s, 2H), 3.11 (d, J=12.6 Hz, 2H), 2.53 (m, 1H), 2.07 (d, J=11.0 Hz, 1H) GCMS m/e 249 (M⁺).

EXAMPLE 12

10 6-METHYL-7-PROPYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,5,8-TETRAENE HYDROCHLORIDE

A) 4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester

- 4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene, (1.9 g, 7.6 mmol) was stirred in 1,4-dioxane (75 mL) and treated with saturated aqueous Na₂CO₃ solution (10 mL).
 15 To this was added di-t-butylidicarbonate (3.31 g, 15.2 mmol). After stirring 6 hours the reaction was treated with H₂O (50 mL) and extracted with EtOAc (4 x 25 mL), dried (Na₂SO₄), filtered, concentrated and chromatographed to provide product (1.9 g, 71%). (TLC 30% EtOAc/hexanes (NH₃) R_f 0.58). ¹H NMR (400 MHz, CDCl₃) δ 7.77 (br s, 1H), 7.72 (br s, 1H), 4.08 (m, 1H), 3.92 (m, 1H), 3.39 (br s, 1H), 3.27 (br s, 1H), 3.25 (m, 1H), 3.18 (m, 1H), 2.46
 20 (m, 1H), 2.02 (d, J=11.0 Hz, 1H).

B) 4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester

- 4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (1.9 g, 5.44 mmol) was hydrogenated in MeOH under a H₂ atmosphere (45 psi) over 10%Pd/C (100 mg) for 1.5 hours then filtered through a Celite pad and concentrated to white solids (1.57 g, 100%). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.14)

- 30 C) 6-Methyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester (For conditions, see: Segelstein, B. E., Chenard, B. L.; Macor, J. E., Post, R. J. *Tetrahedron Lett.* **1993**, 34, 1897)

- 4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (700 mg, 2.42 mmol) was dissolved in EtOH (10 mL) and acetic acid (HOAc) (1 mL) and treated with 1-ethoxyethylenemalononitrile (329 mg, 2.42 mmol). The resulting
 35 mixture was warmed to 60 °C and stirred 18 hours. The reaction was cooled, concentrated treated with H₂O and saturated aqueous Na₂CO₃ solution and extracted with EtOAc (3 x 50 mL), then dried (Na₂SO₄). After filtration and concentration, the residue was

5 chromatographed to provide brown solids (247 mg, 36%) (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.28).

D) 6-Methyl-7-propyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester (For conditions, see; Pilarski, B. *Liebigs Ann. Chem.* **1983**, 1078.)

10 6-Methyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester (80 mg, 0.267 mmol) was stirred in 50% aqueous NaOH solution (3 mL) and DMSO (1 mL) then treated with 1-iodopropane (0.03 mL, 0.321 mmol). This mixture was warmed to 40 °C for 2 hours then cooled, treated with H₂O and extracted with EtOAc. The organic layer was washed with H₂O (3x) then dried (Na₂SO₄), filtered and
15 concentrated to an oil (90 mg, 0.253 mmol). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.15).

E) 6-Methyl-7-propyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene hydrochloride

6-Methyl-7-propyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-
20 tetraene-13-carboxylic acid tert-butyl ester (90 mg, 0.253 mmol) was dissolved in 3N HCl EtOAc (5 mL) and warmed to 100 °C for 1/2 hour. The mixture was cooled, concentrated, slurried in EtOAc, and filtered to provide a white solid (25 mg, 34%). ¹H NMR (400 MHz, DMSO-d₆) δ 9.56 (s, NH), 7.91 (s, 1H), 7.83 (br m, NH), 7.74 (s, 1H), 4.38 (m, 2H), 3.48 (m, 2H), 3.32 (m, 2H), 3.10 (m, 2H), 2.87 (s, 3H), 2.28 (m, 1H), 2.15 (d, J=11.0 Hz, 1H) 1.85 (m,
25 2H), 0.97 (m, 3H). mp 147-150 °C.

EXAMPLE 13

5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8-TETRAENE HYDROCHLORIDE

30 A) 5,7,13-Triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester (For conditions, see; Segelstein, B. E.; Chenard, B. L.; Macor, J. E., Post, R. J. *Tetrahedron Lett.* **1993**, 34, 1897.)

4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (1.0 g, 3.45 mmol) was dissolved in EtOH (10 mL) and HOAc (1 mL) and treated
35 with ethoxymethylenemalononitrile (421 mg, 3.45 mmol). The resulting mixture was warmed to 60 °C and stirred 18 hours. The reaction was cooled, concentrated treated with H₂O and saturated aqueous Na₂CO₃ solution and extracted with EtOAc (3 x 50 mL), then dried

5 (Na₂SO₄). After filtration and concentration, the residue was chromatographed to provide brown solids (580 mg, 56%). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.28)

10 B) 5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene hydrochloride
5,7,13-Triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic
 acid tert-butyl ester was converted to the title compound by the methods described in
 Example 12E. ¹H NMR (400 MHz, D₂O) δ 8.95 (s, 1H), 7.67 (s, 2H), 3.45 (br s, 2H), 3.31 (d,
 J=12.5 Hz, 2H), 3.13 (d, J=12.5 Hz, 2H), 2.30 (m, 1H), 1.99 (d, J=11.5 Hz, 1H). APCI MS m/e
 200.1 [(M + 1)⁺]. mp >250 °C.

15

EXAMPLE 14

7-METHYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8- TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 12D, 5,7,13-
 triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl
 20 ester was converted to the title compound by reaction with iodomethane followed by
 deprotection as described in Example 12E. ¹H NMR (400 MHz, D₂O) δ 8.97 (s, 1H), 7.71 (s,
 1H), 7.67 (s, 1H), 3.94 (s, 3H), 3.48 (m, 2H), 3.33 (d, J=12.2 Hz, 2H), 3.14 (d, J=12.2 Hz, 2H),
 2.34 (m, 1H), 2.03 (d, J=11.5 Hz, 1H). APCI MS m/e 214.2 [(M + 1)⁺].

25

EXAMPLE 15

6-METHYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8- TETRAENE HYDROCHLORIDE

6-Methyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-
 carboxylic acid tert-butyl ester was converted to the title compound by the methods described
 30 in Example 12E. ¹H NMR (400 MHz, DMSO-d₆) δ 9.40 (br m, NH), 7.77 (br m, NH), 7.70 (s,
 1H), 3.44 (m, 2H), 3.30 (m, 2H), 3.05 (br d, J=11.0 Hz, 2H), 2.79 (s, 3H), 2.23 (m, 1H), 2.10
 (d, J=10.8 Hz, 1H). GCMS m/e 213.5 (M⁺).

EXAMPLE 16

35 6,7-DIMETHYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA- 2(10),3,5,8-TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 12D, 6-methyl-5,7,13-
 triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl

5 ester was converted to the title compound by reaction with iodomethane followed by deprotection as described in Example 12E. ¹H NMR (400 MHz, DMSO-d₆) δ 9.52 (s, NH), 7.84 (s, 1H), 7.82 (br m, NH), 7.72 (s, 1H), 3.90 (s, 3H), 3.45 (m, 2H), 3.28 (m, 2H), 3.04 (m, 2H), 2.82 (s, 3H), 2.23 (m, 1H), 2.12 (d, J=11.0 Hz, 1H). APCI MS *m/e* 228.2 [(M + 1)⁺] mp 225-230 °C.

10

EXAMPLE 17

7-PROPYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,5,8-TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 12D, 5,7,13-
 15 triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester was converted to the title compound by reaction with iodopropane followed by deprotection as described in Example 12E. ¹H NMR (400 MHz, DMSO-d₆) δ 9.52 (s, 1H), 9.45 (br s, NH), 7.97 (s, 1H), 7.85 (s, 1H), 7.83 (br m, NH), 4.43 (m, 2H), 3.49 (m, 2H), 3.33 (m, 2H), 3.08 (m, 2H), 2.28 (m, 1H), 2.15 (d, J=11.0 Hz, 1H), 1.92 (m, 2H), 0.93 (m, 3H). APCI
 20 MS *m/e* 242.2 [(M + 1)⁺]. mp 170-171 °C (subl.).

EXAMPLE 18

7-BUTYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,5,8-TETRAENE HYDROCHLORIDE

25 A) 4-Butylamino-5-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (For conditions, see; Senskey, M. D.; Bradshaw, J. D.; Tessier, C. A.; Youngs, W. J. *Tetrahedron Lett.* **1995**, 36, 6217.)

4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (500 mg, 1.43 mmol) and 1-butylamine (1.42 mL, 14.3 mmol) were combined in
 30 THF (5 mL) and stirred 4 hours. The mixture was diluted with EtOAc (50 mL) and washed with H₂O (3 x 30 mL) then dried (Na₂SO₄), filtered and concentrated to an oil. This oil was passed through a Silica gel filter column to remove baseline impurities eluting with 30% EtOAc/hexanes (510 mg, 1.41 mmol, 99%).

35 B) 4-Butylamino-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester

4-Butylamino-5-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (460 mg, 1.27 mmol) was treated with ammonium formate (850 mg, 12.7

5 mmol) and 10%Pd(OH)₂/C (50 mg) in MeOH (20 mL) and brought to reflux for 1 hour then filtered through a Celite pad and concentrated. The solids were treated with saturated aqueous Na₂CO₃ solution, extracted with CH₂Cl₂ (3 x 30 mL) and dried by filtration through a cotton plug to give an oil (440 mg, 100%)

10 C) 7-Butyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester

4-Butylamino-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (440 mg, 1.27 mmol) was dissolved in EtOH (20 mL) and HOAc (2 mL) and treated with ethoxymethylenemalononitrile (186 mg, 1.52 mmol). The resulting mixture
15 was warmed to 60 °C and stirred 18 hours. The reaction was cooled, concentrated, treated with H₂O and saturated aqueous Na₂CO₃ solution then extracted with EtOAc (3 x 50 mL) and dried (Na₂SO₄). After filtration and concentration, the residue was chromatographed to provide a yellow oil (400 mg, 89%) (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.70)

20 D) 7-Butyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene hydrochloride

7-Butyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester was converted to the title compound by the methods described in Example 12E. ¹H NMR (400 MHz, DMSO-d₆) δ 9.93 (brs, NH), 9.68 (s, 1H), 7.99 (s, 1H),
25 7.92 (br m, NH), 7.87 (s, 1H), 4.50 (m, 2H), 3.49 (m, 2H), 3.30 (m, 2H), 3.08 (m, 2H), 2.26 (m, 1H), 2.15 (d, J=11.0 Hz, 1H), 1.88 (m, 2H), 1.32 (m, 2H), 0.82 (t, J=7.0 Hz, 3H) APCI MS *m/e* 256.2 [(M + 1)⁺] mp 204-208 °C

EXAMPLE 19

30 7-Isobutyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene hydrochloride

4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester and isobutylamine were converted to the title compound utilizing the methods described in Example 18A-D. ¹H NMR (400 MHz, CDCl₃) δ 7.74 (s, 1H), 7.52 (s, 1H), 7.14 (s,
35 1H), 3.90 (dd, J=7.5,2.0 Hz, 2H), 3.04-2.97 (m, 4H), 2.70 (dd, J=12.8,2.3 Hz, 2H), 2.42 (m, 1H), 2.19 (m, 1H), 1.98 (d, J=10.5 Hz, 1H), 0.93 (m, 6H) APCI MS *m/e* 256.2 [(M + 1)⁺]. mp 147-150 °C (subl.).

5

EXAMPLE 20

6-METHYL-7-ISOBUTYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8-TETRAENE HYDROCHLORIDE

10 A) 6-Methyl-7-isobutyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester

4-Amino-5-isobutylamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (250 mg, 0.74 mmol) from Example 19B was dissolved in EtOH (10 mL) and HOAc (2 mL) and treated with 1-ethoxyethylenemalononitrile (118 mg, 0.87 mmol). The reaction proceeded as in Example 18C (18h) and was worked up similarly to
15 provide product (TLC 3% MeOH/CH₂Cl₂ (NH₃) R_f 0.57).

B) 6-Methyl-7-isobutyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene hydrochloride

20 6-Methyl-7-isobutyl-5,7,13-triazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,5,8-tetraene-13-carboxylic acid tert-butyl ester was converted to the title compound by the methods described in Example 12E. APCI MS *m/e* 270 3 [(M + 1)⁺] mp 129-130 °C (subl.)

EXAMPLE 21

25 7-PHENYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8-TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 18A, 4,5-dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester and aniline were converted to 4-phenylamino-5-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl at 75 °C for 4 hours in the coupling step. This was then converted to
30 the title compound utilizing the methods described in Example 18B,C,D ¹H NMR (400 MHz, DMSO-d₆) δ 9.08 (1H), 7.78-7.57 (m, 7H), 3.47-3.00 (m, 6H), 2.23 (m, 1H), 2.09 (d, J=11.5 Hz, 1H). APCI MS *m/e* 276.2 [(M + 1)⁺]. mp 210-213 °C.

EXAMPLE 22

35 6-METHYL-7-PHENYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,5,8-TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 21 and Example 20, 4,5-dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester and aniline were

0402010 092859 668260 07020460

5 converted to the title compound. ^1H NMR (400 MHz, DMSO-d_6) δ 7.79 (s, 1H), 7.73-7.56 (m, 5H), 7.32 (s, 1H), 3.46-2.99 (m, 6H), 2.66 (s, 3H), 2.23 (m, 1H), 2.08 (d, $J=11.0$ Hz, 1H). APCI MS m/e 290.2 $[(M + 1)^+]$. mp $>250^\circ\text{C}$.

EXAMPLE 23

10 7-NEOPENTYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,5,8-TETRAENE HYDROCHLORIDE

Utilizing the methods described in Example 18A-D, 4,5-dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester and neopentylamine were converted to the title compound. t-Boc precursor GCMS m/e 369 (M^+).
15 (HCl salt) mp $>250^\circ\text{C}$.

EXAMPLE 24

6-METHYL-7-NEOPENTYL-5,7,13-TRIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,5,8-TETRAENE HYDROCHLORIDE

20 Utilizing the methods described in Example 21 and 20, 4,5-dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester and neopentylamine were converted to the title compound. ^1H NMR (400 MHz, DMSO-d_6) δ 7.31 (s, 1H), 7.27 (s, 1H), 7.02 (br s, 1H), 4.41 (t, $J=13.0$ Hz, 2H), 3.90 (s, 3H), 3.47-3.26 (m, 6H), 2.20 (m, 1H), 2.00 (d, $J=11.5$ Hz, 1H), 0.90 (s, 9H). t-Boc precursor APCI MS m/e 384.2 $[(M + 1)^+]$ mp $>250^\circ\text{C}$.
25

EXAMPLE 25

6,7-DIMETHYL-5,8,14-TRIAZATETRACYCLO[10.3.1.0^{2,11}.0^{4,9}]HEXADEC-2(11),3,5,7,9-PENTAENE HYDROCHLORIDE (Based on the following procedure: Jones, R. G.; McLaughlin, K. C. *Org. Syn.* **1963**, 4, 824. b) Ehrlich, J., Bobert, M. T. *J. Org. Chem.* **1947**, 522.)
30

4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (100 mg, 0.35 mmol) was warmed to 80°C in H_2O (5 mL). To this butane 2,3-dione (0.034 mL, 0.38 mmol) was added under N_2 for 2 hours. The reaction was cooled to room temperature and extracted with EtOAc (3 x 40 mL). The combined organic layer was
35 washed with H_2O (2 x 30 mL), dried (Na_2SO_4), filtered, concentrated and chromatographed on Silica gel to provide an oil (120 mg, 100%). The oil was dissolved in 2N HCl MeOH (5 mL) and warmed to reflux for 30 minutes, then concentrated. Recrystallization from MeOH/ Et_2O provided a white powder (50 mg, 43%) (TLC EtOAc R_f 0.14) ^1H NMR (400 MHz, DMSO-d_6)

00402010 07020460 658260

- 5 δ 7.85 (s, 2H), 3.50 (br s, 2H), 3.32 (d, J=12.5 Hz, 2H), 3.10 (d, J=12.5 Hz, 2H), 2.64 (s, 6H), 2.24 (m, 1H), 2.13 (d, J=11.0 Hz, 1H). t-Boc precursor APCI MS *m/e* 340.3 [(M + 1)⁺].

EXAMPLE 26

5,8,14-TRIAZATETRACYCLO[10.3.1.0^{2,11}.0^{4,9}]HEXADECA-2(11),3,5,7,9-PENTAENE

10 HYDROCHLORIDE

A) 1-(4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone

- 1-(4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (3.0 g, 8.70 mmol) was hydrogenated in MeOH (30 ml) under H₂ (45 psi) over
 15 Pd(OH)₂ (300 mg of 20 wt%/C, 10%wt). After 2.5 hours the reaction was filtered through a Celite pad and rinsed with MeOH (30 ml). The solution was concentrated to a light brown oil which crystallized (2.42 g, 96%). (TLC 10% MeOH/CH₂Cl₂ R_f 0.56). APCI MS *m/e* 286.2 [(M + 1)⁺] mp 129-131 °C.

20 B) 1-(5,8,14-Triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,5,7,9-pentaene)-2,2,2-trifluoro-ethanone

- 1-(4,5-Diamino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (500 mg, 1.75 mmol) was stirred in THF (2 ml). This mixture was treated with H₂O (2 mL) and glyoxal sodium bisulfite addition compound hydrate (931 mg, 3.50 mmol) then
 25 stirred at 55 °C for 2.5 hours. The reaction was cooled to room temperature and extracted with EtOAc (3 x 40 ml). The combined organic layer was washed with H₂O (2 x 30 ml), dried (Na₂SO₄), filtered, concentrated and chromatographed on Silica gel to provide an off white powder (329 mg, 60%) (TLC 25% EtOAc/hexanes R_f 0.40). mp 164-166 °C.

30 C) 5,8,14-Triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,5,7,9-pentaene hydrochloride

- 1-(5,8,14-Triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,5,7,9-pentaene)-2,2,2-trifluoro-ethanone (320 mg, 1.04 mmol) was slurried in MeOH (2.0 ml) and treated with Na₂CO₃ (221 mg, 2.08 mmol) in H₂O (2.0 ml). The mixture was warmed to 70 °C for 2 hours,
 35 then concentrated, treated with H₂O (20 mL) and extracted with CH₂Cl₂ (3 x 10 ml). The organic layer was dried through a cotton plug and concentrated to give a light yellow oil (183 mg, 83%) which solidified upon standing (mp 138-140 °C). This material was dissolved in MeOH (10 mL), treated with 3M HCl/EtOAc (3 ml), concentrated and azeotroped with MeOH

5 (2 x 20 mL) to give solids which were recrystallized from MeOH/Et₂O to afford product as a white solid (208 mg, 97%). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.26). ¹H NMR (400 MHz, CD₃OD) δ 8.94 (s, 2H), 8.12 (s, 2H), 3.70 (m, 2H), 3.54 (d, J=12.5 Hz, 2H), 3.35 (d, J=12.5 Hz, 2H), 2.49 (m, 1H), 2.08 (d, J=11.0 Hz, 1H) GCMS *m/e* 211 (M⁺). mp 225-230 °C.

10

EXAMPLE 27

14-METHYL-5,8,14-TRIAZATETRACYCLO[10.3.1.0^{2,11}.0^{4,9}]HEXADECA-2(11),3,5,7,9-PENTAENE HYDROCHLORIDE

5,8,14-Triazatetracyclo[10.3.1.0^{2,11}.0^{4,9}]hexadeca-2(11),3,5,7,9-pentaene (207 mg, 0.98 mmol) was treated with 37% aqueous formaline solution (1 mL) and formic acid (1 mL) then warmed to 80 °C for 1 hour. The reaction was poured into water, made basic (NaOH, pH ~11) and extracted with EtOAc. The organic layer was dried (Na₂SO₄), concentrated and chromatographed on Silica gel to provide a yellow solid. This was stirred in MeOH (2 mL) and treated with 3N HCl EtOAc (2 mL). After concentration the solids were recrystallized from MeOH/Et₂O to afford product as a white solid (70 mg, 27%). (2% MeOH/CH₂Cl₂ (NH₃) R_f 0.47). ¹H NMR (400 MHz, CDCl₃) δ 8.71 (s, 2H), 7.80 (s, 2H), 3.37 (br s, 2H), 3.03 (m, 2H), 2.47 (m, 2H), 2.32 (m, 1H), 2.18 (br s, 3H), 1.84 (d, J=11.0 Hz, 1H). APCI MS *m/e* 226.2 [(M + 1)⁺]. mp >250 °C.

25

EXAMPLE 28

5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,6,8-TETRAENE HYDROCHLORIDE

A) 2,2,2-Trifluoro-1-(4-hydroxy-5-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone
1-(4,5-Dinitro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (900 mg, 2.61 mmol) and potassium acetate (KOAc) (2.6 g, 26.1 mmol) were dissolved in DMSO (10 mL) and warmed with stirring to 100 °C for 16 hours. The mixture was cooled and diluted with H₂O (50 mL) then extracted with 80% EtOAc/hexanes (6 x 25 mL). The organic layer was washed with H₂O (3 x 20 mL), dried (Na₂SO₄), filtered and concentrated and purified by chromatography to give an oil (575 mg, 70%) (TLC 50% EtOAc/hexanes (NH₃) R_f 0.56)

35

5 B) 2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]-dodeca-2(7),3,5-trien-10-yl)-ethanone

2,2,2-Trifluoro-1-(4-hydroxy-5-nitro-10-aza-tricyclo[6.3.1.0^{2,7}]-dodeca-2(7),3,5-trien-10-yl)-ethanone (575 mg, 1.82 mmol) was hydrogenated in MeOH under a H₂ atmosphere at (45 psi) over 10%Pd/C (80 mg) for 15 hours then filtered through a Celite pad and concentrated
10 to white solids (450 mg, 86%). (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.6). ¹H NMR (400 MHz, CD₃OD) δ 6.67-6.59 (m, 2H), 4.12 (m, 1H), 3.73 (m, 1H), 3.73 (m, 1H), 3.51 (m, 1H), 3.07 (m, 2H), 2.24 (m, 1H), 1.94 (d, J=10.5 Hz, 1H). GCMS m/e 286 (M⁺).

15 C) 2,2,2-Trifluoro-1-(5-oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]-pentadeca-2(10),3,6,8-tetraene)-ethanone (Goldstein, S. W.; Dambek, P. J. *J. Het. Chem.* **1990**, 27, 335)

2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]-dodeca-2(7),3,5-trien-10-yl)-ethanone (150 mg, 0.524 mmol), trimethyl orthoformate (0.19 mL, 1.73 mmol) pyridinium-p-toluenesulfonic acid (PPTS, 18 mg, 0.07 mmol) and xylenes (10 mL) were combined under nitrogen and stirred at 135 °C for 18 hours. The mixture was cooled, treated
20 with H₂O and extracted with EtOAc. The extracts were dried (Na₂SO₄), filtered, concentrated and purified by chromatography to give an oil (110 mg, 71%). (TLC 20% EtOAc/hexanes R_f 0.40)

25 D) 5-Oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]-pentadeca-2(10),3,6,8-tetraene hydrochloride

2,2,2-Trifluoro-1-(5-oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]-pentadeca-2(10),3,6,8-tetraene)-ethanone (110 mg, 0.37 mmol) was stirred in MeOH (5 mL) and treated with Na₂CO₃ (78 mg, 0.74 mmol) in H₂O (2 mL). The stirred mixture was warmed to 80 °C for 2 hours, concentrated to solids, diluted with H₂O and extracted with EtOAc (3 x 40 mL). The product
30 was extracted into aqueous 1N HCl solution (2 x 40 mL) which was washed with EtOAc then neutralized with saturated aqueous Na₂CO₃ solution to pH~10. The product was extracted with EtOAc (3 x 40 mL), dried (Na₂SO₄), concentrated and chromatographed on Silica gel to produce an oil. (TLC 5% MeOH/CH₂Cl₂ (NH₃) R_f 0.19).

The oil was dissolved in MeOH and treated with 3N HCl EtOAc (4 mL) then
35 concentrated, stirred in a minimum of CH₂Cl₂ and saturated with hexanes. After 18 hours, the product was collected by filtration (55 mg, 63%). ¹H NMR (400 MHz, CD₃OD) δ 8.47 (s, 1H), 7.70 (s, 1H), 7.65 (s, 1H), 3.41 (m, 2H), 3.30 (m, 2H), 3.10 (d, J=12.5 Hz, 2H), 2.47 (m, 1H), 2.15 (d, J=11.0 Hz, 1H). APCI MS m/e 201.03 [(M + 1)⁺]

05402010 01020450

5

EXAMPLE 29

6-METHYL-5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,6,8-TETRAENE HYDROCHLORIDE

A) 2,2,2-Trifluoro-1-(6-methyl 5-oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,6,8-tetraene)-ethanone

10 2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone (150 mg, 0.524 mmol), triethyl orthoacetate (0.34 mL, 1.83 mmol), pyridinium-p-toluenesulfonic acid (PPTS, 20 mg, 0.08 mmol) and xylenes (10 mL) were combined under nitrogen and stirred at 135 °C for 18 hours. Workup, isolation and purification as in Example 28C provided the title compound (90 mg, 55%).

15

B) 6-Methyl-5-oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,6,8-tetraene hydrochloride

2,2,2-Trifluoro-1-(6-methyl 5-oxa-7,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2(10),3,6,8-tetraene)-ethanone (90 mg, 0.30 mmol) was stirred in MeOH (5 mL) and treated with Na₂CO₃ (61 mg, 0.58 mmol) in H₂O (2 mL). The stirred mixture was warmed to 80 °C for 2 hours, concentrated to solids, diluted with H₂O and extracted with EtOAc (3 x 40 mL). The solution was dried (Na₂SO₄), concentrated, and chromatographed on Silica gel to produce an oil. (TLC 10% MeOH/CH₂Cl₂ (NH₃) R_f 0.18). ¹H NMR (free base) (400 MHz, CDCl₃) δ 7.40 (s, 1H), 7.26 (s, 1H), 3.05-2.98 (m, 4H), 2.72 (d, J=12.8 Hz, 2H), 2.59 (s, 3H), 2.46 (m, 1H), 1.98 (d, J=10.5 Hz, 1H)

25

The oil was dissolved in MeOH and treated with 3N HCl EtOAc (4 mL) then concentrated, stirred in a minimum of CH₂Cl₂ and saturated with hexanes. After 18 hours, the product was collected by filtration (10 mg, 13%) APCI MS m/e 215.2 [(M + 1)⁺] mp >250 °C.

30

EXAMPLE 30

2-FLUORO-N-(5-HYDROXY-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YL)-BENZAMIDE HYDROCHLORIDE

2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone (150 mg, 0.524 mmol), 2-fluorobenzoyl chloride (0.07 mL, 0.576 mmol), pyridinium-p-toluenesulfonic acid (PPTS, 20 mg, 0.08 mmol), pyridine (0.046 mL, 0.576 mmol) and xylenes (5 mL) were combined under nitrogen and stirred at 135 °C for 18 hours. After 24 hours, additional PPTS (50 mg) was added and the material stirred at 135 °C for an additional 24 hours. Workup as above provided crude product (145 mg, 0.375 mmol) which was

35

09402010 07020460

5 combined with $\text{Na}_2\text{CO}_3(\text{s})$ (80 mg, 0.75 mmol) in MeOH (5 mL) and H_2O (2 mL) and heated to reflux. After 3 hours, the reaction was cooled and diluted with water then extracted with CH_2Cl_2 (4 x 40 mL), dried through a cotton plug then chromatographed to remove baseline impurity (5% MeOH/ CH_2Cl_2 (NH_3)). The crude material was treated with excess 3N HCl EtOAc and concentrated, then dissolved in a minimum of MeOH and the solution was
 10 saturated with Et_2O and stirred. After stirring 4 hours the product was collected by filtration (85 mg, 68%). ^1H NMR (400 MHz, CD_3OD) δ 7.99 (m, 2H), 7.59 (m, 1H), 7.36-7.23 (m, 2H), 6.82 (s, 1H), 2.99 (m, 4H), 2.78 (m, 2H), 2.35 (m, 1H), 1.96 (d, $J=10.5$ Hz, 1H). APCI MS m/e 313 1 [$(\text{M} + 1)^+$]. mp 125-130 °C (subl.).

15

EXAMPLE 31

4-CHLORO-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE HYDROCHLORIDE

A) 1-(4-Chloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone

Copper(I)chloride (CuCl) was prepared as follows: CuSO_4 (4.3 g) and NaCl (1.2 g)
 20 were dissolved in hot H_2O (14 mL) sodium bisulfite (NaHSO_3) (1 g) and sodium hydroxide (NaOH) (690 mg) were dissolved in H_2O (7 mL) and added to the hot acidic solution over 5 minutes. The precipitated white solids were filtered and washed with water.

1-(4-Amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (460 mg, 1.7 mmol) was dissolved in H_2O (2 mL) and concentrated HCl solution (1 mL) then cooled to 0 °C and treated with a solution of sodium nitrite (NaNO_2) (275 mg) in H_2O
 25 (1 mL) dropwise. To the resulting solution was added a CuCl (202 mg, prepared as described above, 2.04 mmol) in concentrated HCl solution (2 mL) over 10 minutes (gas evolution observed). The resulting solution was warmed to 60 °C for 15 minutes, then was cooled to room temperature and extracted with EtOAc (4 x 30 mL). After drying over Na_2SO_4 , the
 30 solution was filtered and concentrated to an oil which was filtered through a Silica pad to remove baseline material eluting with 50% EtOAc/hexanes to give an oil (470 mg, 95%).

B) 4-Chloro-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

1-(4-Chloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (470 mg, 1.62 mmol) and Na_2CO_3 (344 mg, 3.24 mmol) in MeOH (30 mL) and H_2O
 35 (10 mL) were heated to reflux. After 2 hours, the reaction was cooled and diluted with water then extracted with EtOAc (4 x 40 mL), dried (Na_2SO_4), filtered and concentrated to a yellow oil. The crude material was treated with excess 3N HCl EtOAc and concentrated, then

0402010 07020460

5 dissolved in a minimum of CH_2Cl_2 and the solution was saturated with hexanes and stirred. After stirring 4 hours the product was collected by filtration (155 mg, 42%). ^1H NMR (free base) (400 MHz, CDCl_3) δ 7.15 (m, 2H), 7.09 (d, $J=8.0$ Hz, 1H), 3.00-2.94 (m, 4H), 2.68, (m, 2H), 2.38 (m, 1H), 1.92 (d, $J=10.5$ Hz, 1H). ^1H NMR (HCl salt) (400 MHz, $\text{DMSO}-d_6$) δ 7.30-7.20 (m, 3H), 3.30-3.15 (m, 6H), 2.37 (m, 1H), 1.89 (d, $J=11.0$ Hz, 1H). APCI MS m/e 194.1
10 $[(M + 1)^+]$.

EXAMPLE 32

10-AZATRICYCLO[6.3.1.0~2,7~]DODECA-2(7),3,5-TRIEN-4-YL CYANIDE HYDROCHLORIDE

15 A) 1-(4-Iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone

1-(4-Amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (500 mg, 1.85 mmol) was dissolved in H_2O (5 mL) and concentrated H_2SO_4 solution (0.5 mL) then cooled to 0 °C and treated with a solution of sodium nitrite (NaNO_2) (140 mg, 2.04 mmol) in H_2O (2 mL) dropwise. Potassium iodide (460 mg, 2.78 mmol) in 1N H_2SO_4
20 solution (0.5 mL) was added over 10 minutes (reaction becomes dark red). The resulting solution was warmed to room temperature and stirred 18 hours. The reaction was quenched with NaHSO_3 and water (pH 2.5) then extracted with EtOAc (4 x 30 mL). After drying (Na_2SO_4), the solution was filtered and concentrated to a yellow oil which was
25 chromatographed on Silica gel to provide a yellow oil. (260 mg, 37%). (TLC 30% EtOAc/hexanes R_f 0.70). (A 5 g scale performed as above yielded 5 g, 67%).

B) 4-Iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester

30 1-(4-Iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (5 g, 13.1 mmol) and 37% saturated aqueous NH_4OH solution (50 mL) were stirred in MeOH (250 mL) for 2 hours then concentrated and azeotroped with MeOH (2 x 50 mL). The resulting product was stirred in 1,4-dioxane (75 mL) and treated with saturated Na_2CO_3 solution (15 mL). To this was added di-*t*-butyldicarbonate (5.71 g, 26.2 mmol). After stirring
35 18 hours the reaction was treated with H_2O (50 mL) and extracted with CH_2Cl_2 (4 x 30 mL), dried (Na_2SO_4), filtered, concentrated and chromatographed on Silica gel (TLC 20% EtOAc/hexanes) to provide product as an oil (4.9 g, 98%).

5 C) 4-Cyano-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (Utilizing the methods described in House, H. O.; Fischer, W. F. *J. Org. Chem* **1969**, 3626.)

CuCN (108 mg, 1.21 mmol) and NaCN (59 mg, 1.21 mmol) were combined in dry DMF (6 mL) and warmed to 150 °C under N₂. Solution occurs in 20 minutes. To this was
10 added 4-iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (232 mg, 0.6 mmol) in DMF (3.5 mL) and the mixture was stirred for 18 hours at 150 °C. The reaction was cooled and diluted with 50% saturated aqueous NaCl solution and extracted with 50% EtOAc/hexanes (3 x 30 mL). After drying (Na₂SO₄), filtration and concentration the product was isolated by chromatography (86 mg, 50%). (TLC 20% EtOAc/hexanes R_f 0.28).

15

D) 10-Azatricyclo[6.3.1.0~2,7~]dodeca-2(7),3,5-trien-4-yl cyanide hydrochloride

4-Cyano-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester was treated with 3N HCl EtOAc (6 mL) and warmed to reflux for 2 hours, then concentrated, dissolved in a minimum of MeOH which was saturated with Et₂O and stirred 18
20 hours. The product was collected by filtration (49 mg, 73%). ¹H NMR (400 MHz, DMSO-d₆) δ 9.66 (br s, NH), 7.86 (br s, NH), 7.74-7.70 (m, 2H), 7.49 (d, J=7.5 Hz, 1H), 3.33-2.97 (m, 6H), 2.17 (m, 1H), 2.01 (d, J=11.0 Hz, 1H). GCMS m/e 184 (M⁺) mp 268-273 °C.

EXAMPLE 33

25 3-(10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YL)-5-METHYL-1,2,4-OXADIAZOLE HYDROCHLORIDE

4-Cyano-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester (300 mg, 1.1 mmol) was stirred in EtOH (10 mL). To this hydroxyl amine hydrochloride (382 mg, 5.5 mmol) and NaOH (242 mg, 6.05 mmol) were added and the mixture was warmed
30 to reflux. After 45 minutes, the reaction was cooled, diluted with H₂O and extracted with EtOAc. The organic layer was dried (Na₂SO₄) and concentrated to afford a yellow solid (110 mg, 0.35 mmol). This solid was dissolved in pyridine (1 mL) and treated with acetyl chloride (0.03 mL, 0.415 mmol) and warmed to 100°C for 18 hours. The reaction was cooled, treated with H₂O and extracted with EtOAc. The organic extracts were washed with water and
35 saturated aqueous NaCl solution, dried (Na₂SO₄) and concentrated. Chromatography on Silica gel afforded product (50 mg, 0.15 mmol). (25% EtOAc/hexanes R_f 0.18). This product was treated with 2N HCl MeOH (10 mL), heated to 70 °C for 1 hour, cooled, concentrated and recrystallized from MeOH/Et₂O to provide product (15 mg). APCI MS m/e 242.2 [(M + 1)⁺].

0402010 0920460

5

EXAMPLE 34

1-(10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-YL)-1-ETHANONE
HYDROCHLORIDE

A) 1-(4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
ethanone

10 1-(10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (253
mg, 1.0 mmol) and AcCl (0.68 mL, 10 mmol) were dissolved in DCE (3 mL) and treated with
aluminum chloride (AlCl₃) (667 mg, 5.0 mmol). The resulting yellow mixture was stirred for 30
minutes then poured over ice and saturated aqueous NaHCO₃ solution. After stirring 20
minutes the mixture was extracted with CH₂Cl₂ (3 x 30 mL). The organic layer was dried
15 through a cotton plug then concentrated to a orange-yellow oil (255 mg, 86%).

B) 4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-
butyl ester

20 1-(4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
ethanone (1.3 g, 4.37 mmol) and 37% aqueous NH₄OH solution (10 mL) were stirred in MeOH
(30 mL) for 3 hours, then concentrated and azeotroped with MeOH (2 x 50 mL). (This product
could be converted to an HCl salt directly: see the next example.) The resulting product was
stirred in 1,4-dioxane (20 mL) and treated with saturated aqueous Na₂CO₃ solution (5 mL). To
this was added di-*t*-butyldicarbonate (1.91 g, 8.74 mmol). After stirring 2 hours, the reaction
25 was treated with H₂O (50 mL), extracted with CH₂Cl₂ (4 x 30 mL), dried (Na₂SO₄), filtered,
concentrated and chromatographed to provide an oil (1.3 g, 100%). (TLC 40%
EtOAc/hexanes R_f 0.56)

C) 1-(10-Azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-1-ethanone hydrochloride

30 4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl
ester (190 mg, 0.63 mmol) was treated with excess 3N HCl EtOAc and warmed to 70°C for 1
hour then concentrated and dissolved in a minimum of MeOH. The resulting solution was
saturated with Et₂O and stirred. After 18 hours the white crystalline product was collected by
filtration (81 mg, 54%). ¹H NMR (400 MHz, DMSO-d₆) δ 9.75 (br s, NH), 7.89 (s, 1H), 7.88 (d,
J=8.0 Hz, 1H), 7.74 (br s, NH), 7.44 (d, J=8.0 Hz, 1H), 3.33 (br s, 2H), 3.22 (br s, 2H), 3.00 (br
35 m, 2H), 2.54 (s, 3H), 2.17 (m, 1H), 2.02 (d, J=11.0 Hz, 1H). GCMS *m/e* 201 (M⁺). mp 198-202
°C.

05432010 092359
553250 " 07020450

5

EXAMPLE 3510-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIEN-4-OL HYDROCHLORIDE

A) Acetic acid 10-trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl ester

1-(4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-
 10 ethanone (2.5 g, 8.41 mmol) and 3-chloroperoxybenzoic acid (m-CPBA) (7.5 g, 42 mmol) were stirred in CH₂Cl₂ (20 mL) and warmed to 40°C for 18 hours. The mixture was cooled to room temperature, then treated with dimethylsulfide (Me₂S) (3 mL, 40.8 mmol) and stirred 24 hours. The resulting mixture was poured into ice and saturated aqueous Na₂CO₃ solution (100 mL) then extracted with Et₂O (4 x 40 mL). The organic layer was washed saturated
 15 aqueous Na₂CO₃ solution (3 x 40 mL) then dried (Na₂SO₄), filtered and concentrated to afford an oil (1.83 g, 69%). (TLC EtOAc R_f 0.80).

B) 2,2,2-Trifluoro-1-(4-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone

20 Acetic acid 10-trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl ester (900 mg, 2.87 mmol) was stirred in MeOH (20 mL) and saturated aqueous NaHCO₃ solution (15 mL) for 48 hours. The mixture was concentrated, diluted with H₂O and extracted with CH₂Cl₂ (3 x 20 mL) then dried through a cotton plug. Chromatography on Silica gel provided pure product (420 mg, 54%). (TLC 5% MeOH/CH₂Cl₂ R_f 0.44). ¹H NMR (400 MHz, CDCl₃) δ
 25 7.05 (m, 1H), 6.70 (m, 1H), 6.62 (m, 1H), 4.32 (m, 1H), 3.84 (m, 1H), 3.48 (m, 1H), 3.21 (br s, 1H), 3.16 (br s, 1H), 3.09 (m, 1H), 2.38 (m, 1H), 1.97 (d, J=11.0 Hz, 1H)

C) 10-Azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-ol hydrochloride

2,2,2-Trifluoro-1-(4-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-
 30 ethanone (50 mg, 0.184 mmol) was dissolved in MeOH/H₂O (3/1, 5 mL), treated with Na₂CO₃(s) (40 mg, 0.369 mmol) and warmed to 65°C for 2 hours. The mixture was concentrated, diluted with H₂O and extracted with CH₂Cl₂ (3 x 20 mL) then dried through a cotton plug. Filtration through a Silica gel plug provided an oil (10% MeOH/CH₂Cl₂) which was treated with 3N HCl EtOAc (3 mL) then concentrated, dissolved in a minimum of MeOH
 35 which was saturated with Et₂O and stirred. After 18 hours the white crystalline product was collected by filtration (10 mg, 26%). ¹H NMR (400 MHz, CDOD₃) δ 7.16 (d, J=8.0 Hz, 1H), 6.80 (d, J=2.0 Hz, 1H), 6.72 (dd, J=8.0,2.0 Hz, 1H), 3.32-3.28 (4H), 3.09 (dd, J=14.5,12.0 Hz, 2H), 2.32 (m, 1H), 2.03 (d, J=11.0 Hz, 1H) APCI MS m/e 176.2 [(M + 1)⁺] mp 308 (dec.) °C.

558250 "CT020460

EXAMPLE 36

7-METHYL-5-OXA-6,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-
2,4(8),6,9-TETRAENE HYDROCHLORIDE

A) 1-(4-Acetyl-5-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]
trifluoro-ethanone

10 Acetic acid 10-trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl ester
(800 mg, 2.55 mmol) was combined with AlCl₃ (1.0 g, 7.65 mmol) and warmed to 170°C for 2
hours. The mixture was cooled and treated with 1N aqueous HCl solution (20 mL), extracted
with EtOAc and dried (Na₂SO₄). Chromatography affords an oil (190 mg, 24%). (TLC EtOAc
R_f 0.75). ¹H NMR (400 MHz, CDCl₃) δ 12.58 (s, 0.5H), 12.52 (s, 0.5H), 7.53 (s, 1H), 6.86 (s,
15 1H), 4.33 (m, 1H), 3.91 (m, 1H), 3.56 (m, 1H), 3.28 (br s, 1H), 3.24 (br s, 1H), 3.14 (m, 1H),
2.35 (m, 1H), 1.97 (br d, J=11.2 Hz, 1H).

B) 2,2,2-Trifluoro-1-[4-hydroxy-5-(1-hydroxyimino-ethyl)-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl]-ethanone

20 1-(4-Acetyl-5-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7).3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (190 mg, 0.605 mmol), hydroxylamine HCl (99 mg, 1.21 mmol) and NaOAc (118 mg, 1.21 mmol) were combined in MeOH (4 mL) and H₂O (1 mL) and warmed to 65°C for 18 hours. The mixture was cooled, diluted with H₂O and extracted with EtOAc which was dried (Na₂SO₄), filtered and concentrated to provide a yellow oil (177 mg, 93%).

C) 2,2,2-Trifluoro-7-Methyl-5-oxa-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-
2,4(8),6,9-tetraene-ethanone

The above oil, 2,2,2-trifluoro-1-[4-hydroxy-5-(1-hydroxyimino-ethyl)-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl]-ethanone (177 mg, 0.54 mmol) was stirred in DCE (3 mL), treated with triethylamine (0.4 mL, 2.8 mmol) and acetic anhydride (Ac₂O) (0.3 mL, 2.8 mmol) then stirred 18 hours. The reaction was treated with H₂O and extracted with EtOAc. The extracts were dried (Na₂SO₄), filtered and concentrated to a yellow oil which was dissolved in anhydrous DMF (3 mL) and treated with 60% NaH in oil (32 mg, 1.08 mmol). After stirring 18 hours, additional 60% NaH in oil was introduced (33 mg) and the mixture was stirred 2 hours. The reaction was quenched with H₂O (5 mL) and extracted with 80% EtOAc/hexanes (3 x 30 mL). The organic layer was washed with H₂O (3 x 20 mL), dried (Na₂SO₄), filtered and concentrated and chromatographed to provide an oil (40% EtOAc/hexanes R_f 0.56).

5

D) 7-Methyl-5-oxa-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2,4(8),6,9-tetraene hydrochloride

Utilizing the methods described in Example 9C, 2,2,2-Trifluoro-7-Methyl-5-oxa-6,13-diazatetracyclo[9.3.1.0^{2,10}.0^{4,8}]pentadeca-2,4(8),6,9-tetraene-ethanone was converted to the
10 title compound. This was treated with 3N HCl EtOAc (3 mL), concentrated and dissolved in a minimum of CH₂Cl₂ which was saturated with hexanes and stirred. After 18 hours the white crystalline product was collected by filtration (18 mg, 13% overall). ¹H NMR (400 MHz, DMSO-d₆) δ 7.72 (s, 1H), 7.63 (s, 1H), 3.42-2.98 (m, 6H), 2.50 (s, 3H), 2.23 (m, 1H), 2.08 (d, J=10.5 Hz, 1H). APCI MS *m/e* 215.2 [(M + 1)⁺].

15

EXAMPLE 37

4-(2-Methyl-2H-pyrazol-3-yl)-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride and 4-(1-Methyl-1H-pyrazol-3-yl)-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

20 1-(4-Acetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (1.0 g, 3.3 mmol) and dimethylformamide dimethylacetal (DMF-DMA) (4.0 g, 33.6 mmol) were warmed to 140°C for 18 hours. After cooling, a crystalline precipitate was filtered and rinsed with EtOAc (690 mg, 58%).

The above solid, 3-dimethylamino-1-(10-trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-4-yl)-propanone, (200 mg, 0.56 mmol) was dissolved
25 in EtOH (2 mL) and treated with 5N HCl EtOH (0.1 mL) followed by methyl hydrazine (0.6 mmol). The resulting mixture was warmed to 70°C for 4 hours. The mixture was cooled, diluted with water and extracted with EtOAc, dried (Na₂SO₄) and concentrated. Chromatography on Silica gel provided a 3/1 mixture of regioisomeric products (130 mg,
30 68%) (TLC 50% EtOAc/hexanes R_f 0.40)

The above oil (130 mg, 0.388 mmol) and Na₂CO₃(s) (82 mg, 0.775 mmol) were stirred in MeOH (10 mL) and H₂O (5 mL) for 18 hours. After cooling the reaction was diluted with water, extracted with CH₂Cl₂ dried through a cotton plug and concentrated. The product was purified by chromatography on Silica gel and concentrated to an oil. The salt was generated
35 with 2N HCl MeOH, concentrated and recrystallized from MeOH/EtOAc to provide a 3/1 mixture of regioisomeric pyrrazoles (85 mg, 58%). (5% MeOH/CH₂Cl₂ (NH₃) R_f 0.25) TFA-precursor APCI MS *m/e* 336.2 [(M + 1)⁺]

0402010 0920460

5

EXAMPLE 38

4,5-DICHLORO-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE

HYDROCHLORIDE

A) 1-(4,5-Dichloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (Based on Campaigne, E.; Thompson, W. *J. Org. Chem.* **1950**, 72, 629.)

10 1-(10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (539 mg, 2.1 mmol) was stirred in CH₂Cl₂ (5 mL) and treated with ICl₃ (s) (982 mg, 4.21 mmol). The resulting orange solution was stirred 0.5 hours, poured into saturated aqueous NaHSO₃ solution (25 mL), extracted with CH₂Cl₂ (3 x 25 mL), dried through a cotton plug and concentrated to an oil (570 mg, 84%) (TLC 50% EtOAc/hexanes R_f 0.62).

15

B) 4,5-dichloro-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

1-(4,5-Dichloro-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (570 mg, 1.75 mmol) was stirred in MeOH (25 mL) and treated with Na₂CO₃(s) (5 g, 47 mmol) in H₂O (5 mL). The stirred mixture was warmed to 70°C for 4 hours, concentrated to solids, diluted with H₂O and extracted with EtOAc (3 x 40 mL). The product was extracted into 1N aqueous HCl solution (2 x 40 mL) which was washed with EtOAc then neutralized with saturated aqueous Na₂CO₃ solution to pH~10. Product was extracted with CH₂Cl₂ (3 x 40 mL), filtered through a cotton plug and concentrated to an oil (400 mg, 100%)

25 The oil was dissolved in MeOH and treated with 3N HCl EtOAc (4 mL) and concentrated, then dissolved in a minimum of MeOH and which was saturated with Et₂O and stirred 18 hours. The product was collected by filtration (210 mg, 45%) (TLC 50% EtOAc/hexanes (NH₃) R_f 0.08) ¹H NMR (400 MHz, DMSO-d₆) δ 7.58 (s, 2H), 3.33-2.97 (m, 6H), 2.18 (m, 1H), 1.99 (d, J=10.5 Hz, 1H) ¹³C NMR (100 MHz, DMSO-d₆) δ 141.02, 130.60, 126.58, 45.54, 40.55, 38.30 GCMS m/e 227, 229 (M⁺) mp 283-291 °C.

30

EXAMPLE 39

N⁴,N⁴-DIMETHYL-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE-4-SULFONAMIDE

HYDROCHLORIDE

35 A) 10-Trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-sulfonyl chloride

1-(10-Aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-2,2,2-trifluoro-ethanone (530 mg, 2.1 mmol) was added to chlorosulfonic acid (2 mL, 30 mmol) and stirred for 5 minutes

09402010 092239 058250 07020450

- 5 The mixture was quenched with ice, extracted with EtOAc, dried (Na_2SO_4), filtered and concentrated to provide an oil (640 mg, 87%). (TLC 30% EtOAc/hexanes R_f 0.15).

B) N^4,N^4 -Dimethyl-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-sulfonamide hydrochloride

- 10 10-Trifluoroacetyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-sulfonyl chloride (320 mg, 0.9 mmol) was stirred in THF (10 mL) and treated with 40% $\text{Me}_2\text{NH}/\text{H}_2\text{O}$ (1.5 mL). After 10 minutes the mixture was concentrated and chromatographed on Silica gel (TLC 30% EtOAc/hexanes R_f 0.31) to provide an oil (256 mg, 78%). This material was dissolved in MeOH (6 mL) and NH_4OH (2 mL) and stirred 18 hours. The mixture was concentrated and
- 15 azeotroped from MeOH (3x) The resulting oil was dissolved in MeOH and treated with 3N HCl EtOAc (4 mL), concentrated, dissolved in a minimum of MeOH and which was saturated with Et_2O and stirred 18 hours. The product was collected by filtration as a white powder (163 mg, 59%). (TLC 10% MeOH/ CH_2Cl_2 (NH_3) R_f 0.54) ^1H NMR (data, free base) (400 MHz, CDCl_3) δ 7.64 (m, 2H), 7.41 (d, $J=8.0$ Hz, 1H), 3.30 (m, 2H), 3.20 (d, $J=12.5$ Hz, 2H), 3.07 (dd, $J=12.5, 2.2$ Hz, 2H), 2.69 (s, 6H) 2.45, (m, 1H), 2.00 (d, $J=11.0$ Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ 128.43, 124.16, 122.75, 46.67, 46.55, 42.11, 39.44, 37.81 GCMS m/e 266 (M^+) (data HCl salt) ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.68-7.52 (3H), 3.38 (m, 2H), 3.24 (m, 2H), 3.04 (m, 2H), 2.58 (s, 6H), 2.22 (m, 1H), 2.04 (d, $J=11.0$ Hz, 1H) GCMS m/e 266 (M^+). Anal Calcd for $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_2\text{HCl}$: C, 51.56, H, 6.32, N, 9.25 Found C, 51.36; H, 6.09; N, 9.09.
- 20
- 25

EXAMPLE 40

4-(1-PYRROLIDINYL SULFONYL)-10-AZATRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE HYDROCHLORIDE

- The pyrrolidine analogue was prepared from 10-trifluoroacetyl-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-4-sulfonyl chloride (320 mg, 0.9 mmol) as by substituting pyrrolidine in the coupling step described in Example 39B. The TFA product was isolated as an oil (314 mg, 89%) Deprotection and conversion to the salt as in Example 39B affords a white powder (189 mg, 63%). (TLC 10% MeOH/ CH_2Cl_2 (NH_3) R_f 0.60). (TLC 50% EtOAc/hexanes R_f 0.65). ^1H NMR (400 MHz, CDCl_3) δ 7.66 (d, $J=8.0$ Hz, 1H), 7.64 (s, 1H), 7.37 (d, $J=8.0$ Hz, 1H), 3.30-3.15 (m, 8H), 3.00 (m, 2H), 2.39 (m, 1H), 1.98 (d, $J=11.5$ Hz, 1H), 1.72 (m, 4H) ^{13}C NMR (100 MHz CDCl_3) δ 146.91, 144.08, 136.65, 127.90, 124.18, 122.36, 50.43, 47.87, 46.80, 46.63, 42.11, 39.63, 25.10. APCI MS m/e 293 [$(\text{M} + 1)^+$]. (data HCl salt) ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.78 (br s, NH), 8.1 (br s, NH), 7.73 (d, $J=1.5$ Hz, 1H), 7.66
- 30
- 35

5 (dd, J=8.0, 15 Hz, 1H), 7.53 (d, J=8.0 Hz, 1H), 3.39-3.01 (10H), 2.21 (m, 1H), 2.04 (d, J=11.0 Hz, 1H), 1.66 (m, 4H). GCMS *m/e* 292 (*M*⁺). Anal. Calcd. For C₁₃H₁₈N₂O₂HCl.1/2MeOH·C, 54.07, H, 6.47; N, 8.51. Found C, 53.98, H, 6.72, N, 8.12

EXAMPLE 41

10 5,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2,4(8),9-TRIEN-6-ONE
HYDROCHLORIDE (The title compound was prepared following the procedures described in Quallich, G. J.; Morrissey, P. M. *Synthesis* **1993**, 51-53, treating 4,5-dinitro-10-azatricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene-10-carboxylic acid tert-butyl ester as an equivalent to an ortho fluoro phenyl moiety.) ¹H NMR (400 MHz, DMSO-d₆) δ 10.42 (s, NH), 9.88 (br s, NH), 7.52 (br s, 1H), 7.15 (s, 1H), 6.79 (s, 1H), 3.41 (d, J=5.0 Hz, 2H), 3.35-3.13 (m, 4H), 2.93 (m, 2H), 2.12 (m, 1H), 1.95 (d, J=11.5 Hz, 1H). APCI MS *m/e* 215.2 [(*M* + 1)⁺].

EXAMPLE 42

20 6-OXO-5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADEC-2(10),3,6,8-TETRAENE HYDROCHLORIDE (For references, see: Nachman, R. J. *J. Het. Chem.* **1982**, 1545)
 2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone (317 mg, 1.11 mmol) was stirred in THF (10 mL), treated with carbonyldiimidazole (269 mg, 1.66 mmol) and warmed to 60°C for 18 hours. The mixture was concentrated, diluted with CH₂Cl₂ (50 mL) and washed with 1N aqueous HCl solution (3 x 10 mL). The organic layer was dried through a cotton plug, concentrated and chromatographed on Silica gel (50% EtOAc/Hexanes) to provide an oil (130 mg). This material converted to the title compound by the methods described in Example 9C. ¹H NMR (400 MHz, DMSO-d₆) δ 11.78 (s, NH), 9.56 (br s, NH), 7.63 (br s, NH), 7.24 (s, 1H), 7.07 (s, 1H), 3.26 (br s, 2H), 3.16 (br t, J=9.5 Hz, 1H), 2.93 (br s, 1H), 2.18 (m, 1H), 1.97 (d, J=11.0 Hz, 1H). APCI MS *m/e* 217.2 [(*M* + 1)⁺].

EXAMPLE 43

35 3-TRIFLUOROMETHYL-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE
HYDROCHLORIDE (See Grunewald, G. L., Paradkar, V. M.; Pazhenchevsky, B.; Pleiss, M. A., Sall, D. J., Seibel, W. L.; Reitz, T. J. *J. Org. Chem.* **1983**, **48**, 2321-2327. Grunewald, G. L.; Markovich, K. M.; Sall, D. J. *J. Med. Chem.* **1987**, **30**, 2191-2208)

The title compound was prepared by the methods described in Example 1 and 2 starting with 2-fluoro-6-trifluoromethylbromobenzene. ¹H NMR (400 MHz, CD₃OD) δ 7.67-7.50

5 (3H), 3.65 (br s, 1H), 3.49-3.42 (m, 2H), 3.29 (s, 1H), 3.28-3.16 (m, 2H), 2.42 (m, 1H), 2.18 (d, J=11.5 Hz, 1H). APCI MS m/e 228.2 $[(M + 1)^+]$. (HCl salt) mp 275-277 °C. Anal. Calcd. for $C_{12}H_{12}F_3N \cdot HCl \cdot 1/3H_2O$: C, 53.44; H, 5.11; N, 5.19. Found C, 53.73; H, 4.83; N, 5.16.

EXAMPLE 44

10 3-PHENYL-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE HYDROCHLORIDE

A) 5-Fluoro-1,4-dihydro-1,4-methano-naphthalene and 5-iodo-1,4-dihydro-1,4-methano-naphthalene

(Eisch, J. J.; Burlinson, N. E. *J. Amer. Chem. Soc.* **1976**, *98*, 753-761. Paquette, L. A.;
15 Cottrell, D. M.; Snow, R. A. *J. Amer. Chem. Soc.* **1977**, *99*, 3723-3733.)

Magnesium turnings (9.37 g, 385 mmol) were stirred in anhydrous THF (1000 mL) in a flame dried 2L 3 neck round bottom flask equipped with a non-equalizing addition funnel with a N₂ flow adapter, magnetic stirrer and efficient condenser equipped with a N₂ flow adapter. The flask was stirred and warmed to reflux by a removable heating mantle. 2,6-
20 Difluoro-iodobenzene (0.3 g) was added followed by 3N EtMgBr in THF (0.3 mL). The addition funnel was charged with an intimate mixture of cyclopentadiene (24.24 g, 367 mmol) and 2,6-difluoro-iodobenzene (88.0 g, 367 mmol). Small portions (~1 mL) of the intimate mixture were introduced to assist initiation (~4x). After ~15 minutes, the reaction initiated (exotherm, and vapor condensation) and heating was maintained as necessary during the
25 addition of the contents of the addition funnel. The reaction was then maintained at reflux for ~1 hour (no SM by GCMS).

The reaction was cooled to room temperature and quenched with H₂O (200 mL) followed by aqueous 1N HCl solution (200 mL) to dissolve the solids. Product was extracted with hexanes (4 x 150 mL). The combined organic layer was washed with saturated aqueous
30 NaHCO₃ solution (150 mL), dried (Na₂SO₄), filtered through a Silica plug with hexanes rinse and concentrated to an oil (70 g). Chromatography on Silica gel eluting with hexanes provided two lots (9.0 and 21.0 g), which contained primarily 5-iodo-1,4-dihydro-1,4-methano-naphthalene. (TLC hexanes R_f 0.63)

B) 5-Iodo-1,2,3,4-tetrahydro-1,4-methano-naphthalene-2,3-diol

35 5-Iodo-1,4-dihydro-1,4-methano-naphthalene (20 g) and N-methyl morpholine N-oxide (17.61 g, 130 mmol) were stirred in acetone (90 mL) and H₂O (13 mL). To this was added a solution of OsO₄ (0.2 mL, 2.5%wt solution in t-BuOH, 0.02 mmol). After 144 hours, florasil (5 g) and saturated aqueous NaHSO₃ solution (3 mL) were added and stirred for 1/2 hour. The

5 mixture was filtered through a Celite pad and the filtrate concentrated to produce an oil which was purified by chromatography on Silica gel eluting with a gradient of hexanes to 100% EtOAc to provide a yellow solid (13.73 g). APCI MS m/e 301.1 [(M - 1)⁺].

C) 10-Benzyl-3-iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

5-iodo-1,2,3,4-tetrahydro-1,4-methano-naphthalene-2,3-diol (8.33 g, 27.6 mmol) and
 10 Et₃NBnCl (10 mg) were vigorously stirred in dichloroethane (25 mL) and H₂O (75 mL) then treated with sodium periodate (6.17 g, 29.0 mmol). After 1.5 hours, the layers were separated and the aqueous layer extracted with DCE (2 x 40 mL). The combined organic layer was washed with H₂O (4 x 30 mL) until no reaction to starch iodide paper was observed, then with saturated aqueous NaCl solution (30 mL). The organic layer was dried through a cotton plug
 15 and treated with benzyl amine (3.16 mL, 29.0 mmol) and stirred for 2 minutes then transferred to an addition funnel. This solution was added over ~10 minutes to a vigorously stirred cooled (0 °C) mixture of NaBH(OAc)₃ (18.72 g, 88.0 mmol) in DCE (150 mL). After addition was complete, the mixture was stirred without cooling for 2 hours. The mixture was quenched with saturated aqueous Na₂CO₃ solution (100 mL) and stirred for 1 hour, then the layers were
 20 separated and the aqueous layer was extracted with CH₂Cl₂ (3 x 50 mL). The combined organic layer was washed with saturated aqueous NaCl solution (50 mL), dried through a cotton plug and concentrated. Chromatography on Silica gel provided an oil (6.3 g, 61%) (TLC 5% EtOAc/hexanes R_f 0.10). ¹H NMR (400 MHz, CDCl₃) δ 7.61 (d, J= 8.0 Hz, 1H), 7.28-7.22 (m, 3H), 7.13 (d, J=8.0 Hz, 1H), 6.98-6.94 (m, 3H), 3.58 (AB dd, J=14.2 Hz, 2H), 3.26 (br s, 1H), 3.21 (br s, 1H), 3.04 (br d, J=10.2 Hz, 1H), 2.83 (br d, J=10.2 Hz, 1H), 2.47 (d, J=10.0 Hz, 1H), 2.39 (d, J=10.0 Hz, 1H), 2.34 (m, 1H), 1.72 (d, J=10.5 Hz, 1H) APCI MS m/e 376.0 [(M + 1)⁺]

D) 10-Benzyl-3-phenyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

(For a discussion, see: Miyaura, N; Suzuki, A *Chem Rev* **1995**, 95, 2457-
 30 2483.)

10-Benzyl-3-iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (375.3 mg, 1.0 mmol), potassium acetate (785 mg, 8.0 mmol) and phenyl boronic acid (183 mg, 1.5 mmol) were combined in 10/1 EtOH/H₂O (5 mL). The mixture was degassed (3 vacuum/N₂ cycles), treated with tetrakis(triphenylphosphine)palladium(0) (57.5 mg, 0.05 mmol) and warmed to 90
 35 °C for 18h. The reaction was cooled, diluted with H₂O and extracted with Et₂O (3 x 50 mL). The organic layer was washed with brine (50 mL), dried (MgSO₄), filtered and concentrated to provide an oil (180 mg, 55%) (TLC 4%EtOAc/hexanes R_f 0.18) GCMS m/e 325 (M)⁺.

E) 3-Phenyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride

5 10-Benzyl-3-phenyl-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene was converted into the title compound utilizing the conditions described in Example 2D (TLC 10% MeOH/CH₂Cl₂ (NH₃) R_f 0.30). (data for free base) ¹H NMR (400 MHz, CDCl₃) δ 7.46-7.15 (8H), 3.17 (br s, 1H), 3.01 (m, 2H), 2.93 (d, J=13.0 Hz, 1H), 2.72 (dd, J=10.5,2.5 Hz, 1H), 2.63 (dd, J=10.5,2.5 Hz, 1H), 2.41 (m, 1H), 1.91 (d, J=10.5 Hz, 1H). APCI MS *m/e* 236.2 [(M + 1)⁺]
10 (HCl salt) mp 262-265 °C. Anal. Calcd. for C₁₇H₁₇N.HCl.1/3H₂O: C, 73.26; H, 6.86; N, 5.19 Found C, 73.50; H, 6.77; N, 5.04.

EXAMPLE 45

3-HYDROXY-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE

15 HYDROCHLORIDE

A) 10-Benzyl-3-boronic acid-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

10-Benzyl-3-iodo-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (3.0 g, 7.99 mmol) was stirred in anhydrous THF (40 mL) at -78 °C under nitrogen and treated dropwise with n-BuLi (3.84 mL of 2.5M soln. in hexanes, 9.59 mmol) After 10 minutes, tri-isopropylborate
20 (4.61 mL, 20.0 mmol) was added dropwise. After ~1/2 hour, the reaction was poured into saturated aqueous NaHCO₃ solution, stirred 5 minutes and extracted with EtOAc (3 x 50 mL) and concentrated The residue was dissolved in 30% Et₂O/hexanes and extracted with 1N NaOH aqueous solution (4 x 50 mL). The combined aqueous basic layer was treated with concentrated HCl to achieve pH 8 and extracted with EtOAc (4 x 25 mL), dried (Na₂SO₄) and
25 stripped. Chromatography on Silica gel eluting first with 3% EtOAc/hexanes to remove non-polar components, then with 5% MeOH/CH₂Cl₂ provides the title compound. (TLC 25% EtOAc/hexanes R_f 0.60)

B) 10-Benzyl-3-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene

10-Benzyl-3-boronic acid-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (140 mg, 0.48 mmol) dissolved in THF (5 mL) was treated with N-methylmorpholine-N-oxide (64.5 mg, 0.48 mmol) and brought to reflux for 1 hour. The reaction was concentrated and chromatographed on Silica gel to provide product. (TLC 25% EtOAc/hexanes R_f 0.18). ¹H
30 NMR (400 MHz, CDCl₃) δ 7.18-7.15 (3H), 7.04 (dd, J= 8.0,7.0 Hz, 1H), 6.95 (m, 2H), 6.75 (d, J=7.0 Hz, 1H), 6.59 (dd, J=8.0,1.0 Hz, 1H), 3.53 (br s, OH), 3.51 (AB d, J=14.0 Hz, 2H), 3.28 (br s, 1H), 3.06 (br s, 1H), 2.91 (dd, J=8.5,1.5 Hz, 1H), 2.79 (ddd, J=8.5,1.5,1.5 Hz, 1H), 2.42 (d, J=11.0 Hz, 1H), 2.39 (d, J=11.0 Hz, 1H), 2.23 (m, 1H), 1.65 (d, J=10.5 Hz, 1H). APCI MS *m/e* 266.5 [(M + 1)⁺]

5 C) 3-Hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene hydrochloride
 10-Benzyl-3-hydroxy-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-triene (160 mg, 0.60 mmol) was converted into the title compound by the methods described in Example 1D. ¹H NMR (400 MHz, CDCl₃) δ 7.15 (dd, J=8.0,7.5 Hz, 1H), 6.84 (d, J=7.5 Hz, 1H), 6.76 (d, J=8.0 Hz, 1H), 3.51 (br s, 1H), 3.33-3.25 (3H), 3.16 (d, J=12.0 Hz, 1H), 3.09 (d, J=12.0 Hz, 1H), 2.29 (m, 1H), 2.02 (d, J=11.0 Hz, 1H). APCI MS *m/e* 175.8 [(M + 1)⁺]. (HCl salt) mp 253-255 °C.

EXAMPLE 46

4,5-DIFLUORO-10-AZA-TRICYCLO[6.3.1.0^{2,7}]DODECA-2(7),3,5-TRIENE HYDROCHLORIDE

The title compound was prepared by the methods described in Example 1 and 2 starting with 2,4,5-trifluorobromobenzene ¹H NMR (400 MHz, CDCl₃) δ 7.31 (t, J=8.5 Hz, 2H), 3.48-3.13 (6H), 2.38 (m, 1H), 2.11 (d, J=11.5 Hz, 1H). APCI MS *m/e* 196.2 [(M + 1)⁺]. (HCl salt) mp 301-303 °C. Anal. Calcd. for C₁₁H₁₁F₂N.HCl.1/6H₂O: C, 56.30; H, 5.30; N, 5.97 Found C, 56.66; H, 5.41; N, 5.96

EXAMPLE 47

6-ETHYL-5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,6,8-TETRAENE HYDROCHLORIDE

2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone and propionyl chloride were converted to the title compound following the procedures described in Example 30 and Goldstein, S. W.; Dambek, P. J. *J. Het. Chem.* **1990**, 27, 335. ¹H NMR (400 MHz, CD₃OD) δ 7.64 (s, 1H), 7.62 (s, 1H), 3.48 (d, J=2.5 Hz, 2H), 3.41 (d, J=12.0 Hz, 2H), 3.20 (2H), 3.01 (q, J=7.5 Hz, 2H), 2.45 (m, 1H), 2.17 (d, J=11.5 Hz, 1H), 1.42 (t, J=7.5 Hz, 3H) APCI MS *m/e* 229.2 [(M + 1)⁺]

EXAMPLE 48

6-ISOPROPYL-5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECA-2(10),3,6,8-TETRAENE HYDROCHLORIDE

2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone and isobutyryl chloride were converted to the title compound following the procedures described in EXAMPLE 47 (TLC 25% EtOAc/hexanes R_f 0.14) ¹H NMR (400 MHz, CD₃OD) δ 7.65 (2H), 3.49 (br s, 2H), 3.41 (d, J=12.0 Hz, 2H), 3.33-3.19 (3H), 2.45 (m, 1H), 2.18 (d, J=11.5 Hz, 1H), 1.45 (d, J=7.0 Hz, 6H). APCI MS *m/e* 243.2 [(M + 1)⁺] (HCl salt) mp 249-251 °C.

5

EXAMPLE 49

6-BENZYL-5-OXA-7,13-DIAZATETRACYCLO[9.3.1.0^{2,10}.0^{4,8}]PENTADECAL-2(10),3,6,8-TETRAENE HYDROCHLORIDE


10 2,2,2-Trifluoro-1-(4-hydroxy-5-amino-10-aza-tricyclo[6.3.1.0^{2,7}]dodeca-2(7),3,5-trien-10-yl)-ethanone and phenyl-acetyl chloride were converted to the title compound following the procedures described in EXAMPLE 47. ¹H NMR (400 MHz, CD₃OD) δ 7.63 (s, 1H), 7.58 (s, 1H), 7.36-7.24 (5H), 4.29 (s, 2H), 3.46 (d, J=2.5 Hz, 2H), 3.39 (d, J=12.0 Hz, 2H), 3.18 (2H), 2.42 (m, 1H), 2.15 (d, J=11.5 Hz, 1H). APCI MS *m/e* 291.2 [(M + 1)⁺].

1563250 " OF 20450

5

CLAIMS

1. A compound of the formula



R^1 is hydrogen, (C_1-C_6) alkyl, unconjugated (C_3-C_6) alkenyl, $XC(=O)R^{13}$ or $-CH_2CH_2-O-(C_1-C_4)$ alkyl;

10 R^2 and R^3 are selected, independently, from hydrogen, (C_2-C_6) alkenyl, (C_2-C_6) alkynyl, hydroxy, nitro, amino, halo, cyano, $-SO_q(C_1-C_6)$ alkyl wherein q is zero, one or two, (C_1-C_6) alkylamino-, $[(C_1-C_6)alkyl]_2$ amino-, $-CO_2R^4$, $-CONR^5R^6$, $-SO_2NR^7R^8$, $-C(=O)R^{13}$, $-XC(=O)R^{13}$, aryl- (C_0-C_3) alkyl- or aryl- (C_0-C_3) alkyl-O-, wherein said aryl is selected from phenyl and naphthyl, heteroaryl- (C_0-C_3) alkyl- or heteroaryl- (C_0-C_3) alkyl-O-, wherein said heteroaryl is
15 selected from five to seven membered aromatic rings containing from one to four heteroatoms selected from oxygen, nitrogen and sulfur, and $X^2(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$, wherein X^2 is absent or X^2 is (C_1-C_6) alkylamino- or $[(C_1-C_6)alkyl]_2$ amino-, and wherein the $(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$ moiety of said $X^2(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$ contains at least one carbon atom, and wherein from one to three of the carbon atoms of said $(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$ moiety may
20 optionally be replaced by an oxygen, nitrogen or sulfur atom, with the proviso that any two such heteroatoms must be separated by at least two carbon atoms, and wherein any of the alkyl moieties of said $(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$ may be optionally substituted with from two to seven fluorine atoms, and wherein one of the carbon atoms of each of the alkyl moieties of said aryl- (C_0-C_3) alkyl- and said heteroaryl- (C_0-C_3) alkyl- may optionally be replaced by an oxygen, nitrogen
25 or sulfur atom and wherein each of the foregoing aryl and heteroaryl groups may optionally be substituted with one or more substituents, preferably from zero to two substituents, independently selected from (C_1-C_6) alkyl optionally substituted with from one to seven fluorine atoms, $(C_1-C_6)alkoxy$ optionally substituted with from two to seven fluorine atoms, halo (e.g., chloro, fluoro, bromo or iodo), (C_2-C_6) alkenyl, (C_2-C_6) alkynyl, hydroxy, nitro, cyano, amino, (C_1-C_6) alkylamino-, $[(C_1-C_6)alkyl]_2$ amino-, $-CO_2R^4$, $-CONR^5R^6$, $-SO_2NR^7R^8$, $-C(=O)R^{13}$ and $-XC(=O)R^{13}$,
30 $X^2(C_0-C_6)alkoxy-(C_0-C_6)alkyl-$.

or R^2 and R^3 , together with the carbons to which they are attached, form a four to seven membered monocyclic or ten to fourteen membered bicyclic, carbocyclic ring that can be saturated or unsaturated, wherein from one to three of the nonfused carbon atoms of said
35 monocyclic rings, and from one to five of the carbon atoms of said bicyclic rings that are not part


- 5 of the benzo ring shown in formula I, may optionally and independently be replaced by a nitrogen, oxygen or sulfur, and wherein said monocyclic and bicyclic rings may optionally be substituted with one or more substituents, preferably from zero to two substituents for the monocyclic rings and from zero to three substituents for the bicyclic rings, that are selected, independently, from (C₁-C₆) alkyl optionally substituted with from one to seven fluorine atoms, (C₁-C₆) alkoxy optionally substituted with from one to seven fluorine atoms, nitro, cyano, halo, (C₂-C₆) alkenyl, (C₂-C₆) alkynyl, hydroxy, amino, (C₁-C₆) alkylamino and [(C₁-C₆) alkyl]₂amino, -CO₂R⁴, -CONR⁵R⁶, -SO₂NR⁷R⁸, -C(=O)R¹³ and -XC(=O)R¹³;

each R⁴, R⁵, R⁶, R⁷, R⁸ and R¹³ is selected, independently, from hydrogen and (C₁-C₆) alkyl, or R⁵ and R⁶, or R⁷ and R⁸ together with the nitrogen to which they are attached, form a pyrrolidine, piperidine, morpholine, azetidine, piperazine, N-(C₁-C₆) alkylpiperazine or thiomorpholine ring, or a thiomorpholine ring wherein the ring sulfur is replaced with a sulfoxide or sulfone; and

each X is, independently, (C₁-C₆) alkylene;

- with the proviso that: (a) at least one of R¹, R² and R³ must be the other than hydrogen, and (b) when R² and R³ are both hydrogen, R¹ cannot be hydrogen or methyl; or a pharmaceutically acceptable salt thereof;

2. A compound according to claim 1, wherein R² and R³, together with the benzo ring of formula I, form a bicyclic ring system selected from the following:



- 25 wherein R¹⁰ and R¹⁷ are selected, independently, from (C₀-C₆) alkoxy-(C₀-C₆) alkyl- wherein the total number of carbon atoms does not exceed six and wherein any of the alkyl moieties may optionally be substituted with from one to seven fluorine atoms, nitro, cyano, halo,

5 amino, (C₁-C₆)alkylamino-, [(C₁-C₆) alkyl]₂amino-, -CO₂R⁴, -CONR⁵R⁶, -SO₂NR⁷R⁸, -C(=O)R¹³,
-XC(=O)R¹³, phenyl and monocyclic heteroaryl, wherein said heteroaryl is selected from five to
seven membered aromatic rings containing from one to four heteroatoms selected from oxygen,
nitrogen and sulfur,

3. A compound according to claim 1, wherein R² and R³ do not, together with the
10 benzo ring of formula I, form a bicyclic or tricyclic ring system.

4. A compound according to claim 1, wherein one or both of R² and R³ are
-C(=O)R¹³ wherein R¹³ is (C₁-C₆)alkyl.

5. A compound according to claim 1, wherein one of R² and R³ is -COR¹³ wherein
R¹³ is (C₁-C₆)alkyl or (C₁-C₃)alkyl optionally substituted with from one to seven fluorine atoms.

15 6. A compound according to claim 1, wherein one of R² and R³ is CF₃, fluoro,
cyano or C₂F₅.

7 A pharmaceutical composition for use in reducing nicotine addiction or aiding in
the cessation or lessening of tobacco use in a mammal, comprising an amount of a compound
according to claim 1 that is effective in reducing nicotine addiction or aiding in the cessation or
20 lessening of tobacco use and a pharmaceutically acceptable carrier.

8. A method for reducing nicotine addiction or aiding in the cessation or lessening
of tobacco use in a mammal, comprising administering to said mammal an amount of a
compound according to claim 1 that is effective in reducing nicotine addiction or aiding in the
cessation or lessening of tobacco use.


25 9. A pharmaceutical composition for treating a disorder or condition selected from
inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma
gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain,
acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar
disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS), cognitive
30 dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid
hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical
dependencies and addictions (e.g. dependencies on, or addictions to nicotine (and/or tobacco
products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke,
traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia,
35 dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including
petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease
(PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal,

5 comprising an amount of a compound according to claim 1 that is effective in treating such disorder or condition and a pharmaceutically acceptable carrier.

10 A method for treating a disorder or condition selected from inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS), cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia, dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal, comprising administering


20 to a mammal in need of such treatment an amount of a compound according to claim 1 that is effective in treating such disorder or condition.

11 A compound of the formula




wherein P is hydrogen, methyl, COOR¹⁶ wherein R¹⁶ is (C₁-C₆)alkyl, allyl or 2,2,2-trichloroethyl, -C(=O)NR⁵R⁶ wherein R⁵ and R⁶ are defined as in formula I above; -C(=O)H, -C(=O)(C₁-C₆)alkyl wherein the alkyl moiety may optionally be substituted with from 1 to 3 halo atoms, preferably with from 1 to 3 fluoro or chloro atoms; benzyl, t-butoxycarbonyl (t-Boc) or trifluoroacetyl, and R¹⁴ and R¹⁵ are selected, independently, from hydrogen, (C₁-C₆)alkyl optionally substituted with from one to seven fluorine atoms; -C(=O)(C₁-C₆)alkyl, cyano, hydroxy, nitro, amino, -O(C₁-C₆)alkyl and halo, with the proviso that R¹⁴ and R¹⁵ can not both be hydrogen when P is hydrogen or methyl

- 5 12. A method for reducing nicotine addiction or aiding in the cessation or lessening of tobacco use in a mammal, comprising administering to said mammal an amount of a compound comprising an amount of a compound of the formula




10 or a pharmaceutically acceptable salt thereof, that is effective in reducing nicotine addiction or aiding in the cessation or lessening of tobacco use.

13. A method for treating a disorder or condition selected from inflammatory bowel disease (including but not limited to ulcerative colitis, pyoderma gangrenosum and Crohn's disease), irritable bowel syndrome, spastic dystonia, chronic pain, acute pain, celiac sprue, pouchitis, vasoconstriction, anxiety, panic disorder, depression, bipolar disorder, autism, sleep disorders, jet lag, amyotrophic lateral sclerosis (ALS), cognitive dysfunction, hypertension, bulimia, anorexia, obesity, cardiac arrhythmias, gastric acid hypersecretion, ulcers, pheochromocytoma, progressive supramuscular palsy, chemical dependencies and addictions (e.g., dependencies on, or addictions to nicotine (and/or tobacco products), alcohol, benzodiazepines, barbituates, opioids or cocaine), headache, stroke, traumatic brain injury (TBI), psychosis, Huntington's Chorea, tardive dyskinesia, hyperkinesia, dyslexia, schizophrenia, multi-infarct dementia, age related cognitive decline, epilepsy, including petit mal absence epilepsy, senile dementia of the Alzheimer's type (AD), Parkinson's disease (PD), attention deficit hyperactivity disorder (ADHD) and Tourette's Syndrome in a mammal, comprising administering to a mammal in need of such treatment an amount of a compound of the formula



25 or a pharmaceutically acceptable salt thereof, that is effective in treating such disorder or condition

14. A compound of the formula



30 wherein R^2 and R^3 are defined as in claim 1, and P' is COOR^{16} wherein R^{16} is allyl, 2,2,2-trichloroethyl or $(\text{C}_1\text{-C}_6)\text{alkyl}$, $-\text{C}(=\text{O})\text{NR}^5\text{R}^6$ wherein R^5 and R^6 are defined as in claim 2.

- 5 -C(=O)H, -C(=O)(C₁-C₆)alkyl wherein the alkyl moiety may optionally be substituted with from 1 to 3 halo atoms, preferably with from 1 to 3 fluoro or chloro atoms; benzyl, or t-butoxycarbonyl (t-Boc).


09402010 092399
1615260 " 01020460

5

ARYL FUSED AZAPOLYCYCLIC COMPOUNDS

Abstract

Compounds of the formula



and their pharmaceutically acceptable salts, wherein R¹, R², R³ and n are defined as in the specification, intermediates in the synthesis of such compounds, pharmaceutical compositions containing such compounds and methods of using such compounds in the treatment of neurological and psychological disorders are claimed.

09402010.0000

Please type a plus sign (+) inside this box →

+

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63) <input checked="" type="checkbox"/> Declaration submitted with Initial Filing <input type="checkbox"/> Declaration Submitted after Initial Filing (surcharge 37 CFR 1.16 (e)) required)		Attorney Docket Number	PC 10030A
		First Named Inventor	Jotham Wadsworth COE
		COMPLETE IF KNOWN	
		Application Number	Not yet assigned
		Filing Date	Filed herewith
		Group Art Unit	Not yet assigned
		Examiner Name	Not yet assigned

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

ARYL FUSED AZAPOLYCYCLIC COMPOUNDS

(Title of the Invention)

the specification of which
☐ is attached hereto

OR

☒ was filed on (MM/DD/YYYY) 11/13/1998 as United States Application Number or PCT International

Application Number PCT/IB98/01813 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
			<input type="checkbox"/>	YES <input type="checkbox"/>	NO <input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below:

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B sheet attached hereto.
60/070,245	12/31/1997	

EXPRESS MAIL NO. EM484852791

Please type a plus sign (+) inside this box →

+

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION ---- Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 U.S.C. 1.56, which became available between the filing date of the prior application and the national or PCT International filing date of this application.

U.S. Parent Application Number or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT International application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent

and Trademark Office connected therewith:

☐ Customer Number

or

Place Customer
Number Bar Code
Label here

☒ Registered practitioner(s) name/registration number listed below

Name	Registration Number	Name	Registration Number
Peter C. Richardson	27,526	Mark Dryer	28,775
Allen J. Spiegel	25,749	Lawrence C. Akers	28,587
Paul H. Ginsburg	28,718	A. Dean Olson	31,185
J. Trevor Lumb	28,567	Mervin E. Brokke	32,723
James T. Jones	30,561	Valerie M. Fedowich	33,688
Gregg C. Benson	30,977	Bryan C. Zielinski	34,462
Robert F. Sheyka	31,304	Robert T. Ronau	36,257
Grover F. Fuller Jr.	31,760	B. Timothy Creagan	39,156
Karen DeBenedictis	32,977	Alan L. Koller	37,371
Lorraine B. Ling	35,251	Jolene W. Appleman	35,428
Garth Butterfield	36,997	Kristina L. Konstas	37,864
Carl J. Goddard	39,203	Seth H. Jacobs	32,140
Raymond M. Speer	26,810	Martha A. Gammill	31,820
Jennifer A. Kispert	40,049	Gregory P. Raymer	36,647
Jacob M. Levine	32,509	E. Victor Donahue	35,492
Israel Nissenbaum	27,582	Roy F. Waldron	42,208
Steven W. Collier	42,429	Todd M. Chrissey	37,807

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.

Direct all correspondence to:

☐ Customer Number
or Bar Code Label

OR ☒ Correspondence address below

Name	<u>Paul H. Ginsburg</u>				
Address	<u>Pfizer Inc.</u>				
Address	<u>235 East 42nd Street, 20th Floor</u>				
City	<u>New York</u>	State	<u>New York</u>	Zip Code	<u>10017-5755</u>
Country	<u>United States Of America</u>	Telephone	<u>(212)573-2369</u>	Fax	<u>(212)573-1939</u>

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: ☐ A petition has been filed for this unsigned inventor


Given Name (first and middle (if any))				Family Name or Surname			
<u>Jotham Wadsworth</u>				<u>Wadsworth Jr</u>			
Inventor's Signature				Date			
<u>Jotham W. Coc</u>				<u>9/22/99</u>			
Residence: City	<u>Niantic</u>	State	<u>CT</u>	Country	<u>US</u>	Citizenship	<u>US</u>
Post Office Address							
<u>8 Bush Hill Drive</u>							
Post Office Address							
City	<u>Niantic</u>	State	<u>CT</u>	Zip	<u>06357</u>	Country	<u>US</u>

☒ Additional inventors are being named on the supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto.

Please type a plus sign (+) inside this box →

+

DECLARATION	ADDITIONAL INVENTOR(S) Supplemental Sheet
--------------------	--

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Paige Roanne Palmer				BROOKS			
Inventor's Signature				Date		9/28/99	
Residence: City	North Stonington	State	CT	Country	US	Citizenship	US
Post Office Address	9 Wyassup Road						
Post Office Address							
City	North Stonington	State	CT	Zip	06359	Country	US

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Inventor's Signature				Date			
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Inventor's Signature				Date			
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	

Name of Additional Joint Inventor, if any:		<input type="checkbox"/> A petition has been filed for this unsigned inventor					
Given Name (first and middle [if any])				Family Name or Surname			
Inventor's Signature				Date			
Residence: City		State		Country		Citizenship	
Post Office Address							
Post Office Address							
City		State		Zip		Country	

EXPRESS MAIL NO. EM484852791